For Research Use Only

STMN2 Polyclonal antibody

Catalog Number: 10586-1-AP

Featured Product

29 Publications



Basic Information

Catalog Number:

10586-1-AP

BC006302

Concentration:

600 ug/ml

11075

Source:

Rabbit

UNIPROT ID:

Rabbit

Q93045

Isotype:

IgG

GenBank Accession Number:

GenBank Accession Number:

BC006302

UNCBI):

UNIPROT ID:

Q93045

Full Name:

stathmin-like 2

Immunogen Catalog Number: Calculated MW: AG0929 21 kDa

Observed MW: 18-21 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications:
WB, IHC, IF
Species Specificity:
human, mouse, rat
Cited Species:

human, mouse, rat, chicken, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:200-1:800

Purification Method:

Background Information

Stathmin-2 (encoded by STMN2), a member of the stathmin family of proteins, encodes a phosphoprotein that has an important role in microtubule dynamics. The STMN2 protein promotes microtubule dynamics in axonal growth cones, and is thought to play an important role in neurite outgrowth. STMN2 protein is localized in both Golgi and cytosolic compartments. Its expression appears modulated in osteoblasts by nerve growth factor, dexamethasone or RhoA kinase inhibitor Y-27632 which are known effectors of osteogenesis. (PMID: 33841129, PMID: 18611392)

Positive Controls:

IP: mouse brain tissue.

IHC: mouse brain tissue,

IF/ICC: SH-SY5Y cells,

WB: mouse brain tissue, SH-SY5Y cells

Notable Publications

Author	Pubmed ID	Journal	Application
Konstantin Feinberg	28877995	J Cell Biol	WB
Adam Tuttle	31476133	Elife	IHC
Andrea Loreto	31740269	Neurobiol Dis	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10586-1-AP (STMN2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 10586-1-AP (STMN2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of STMN2 antibody (10586-1-AP; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-STMN2 transfected SH-SY5Y cells.



IP result of anti-STMN2 (IP:10586-1-AP, 4ug; Detection:10586-1-AP 1:10000) with mouse brain tissue lysate 1840 ug.



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using STMN2 antibody (10586-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).