

For Research Use Only

# BAG3 Polyclonal antibody

Catalog Number: 10599-1-AP

Featured Product

123 Publications



## Basic Information

### Catalog Number:

10599-1-AP

### Concentration:

480 µg/ml

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG0956

### GenBank Accession Number:

BC006418

### GeneID (NCBI):

9531

### UNIPROT ID:

O95817

### Full Name:

BCL2-associated athanogene 3

### Calculated MW:

61 kDa

### Observed MW:

74-80 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:30000-1:60000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:500-1:2000

IF/ICC 1:50-1:500

## Applications

### Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

### Cited Applications:

WB, IHC, IF, IP, CoIP

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse, rat, monkey, hamster

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB:** HEK-293 cells, HeLa cells, K-562 cells, mouse heart, rat heart

**IP:** K-562 cells,

**IHC:** human lung cancer tissue, human gliomas tissue

**IF/ICC:** A549 cells, HeLa cells, HepG2 cells

## Background Information

BAG3 (Bcl2-associated athanogene 3) belongs to the BAG protein family, the co-chaperone that binds to Hsc70/Hsp70 through the BAG domain and modulates their activity in polypeptide folding. BAG3 contains also a WW domain and a proline-rich (PXXP) repeat, that mediate binding to partners different from Hsp70. Through interacting with different molecular partner, BAG3 influences several cell processes, such as apoptosis, autophagy and cell motility. BAG3 protein has been reported to sustain cell survival, resistance to therapy, and/or motility and metastatization in several tumor types, thus being identified as a potential target for anticancer therapies. In addition, defects in BAG3 are the cause of some myopathy. BAG3 normally migrates around 74-80 kDa; a slightly different molecular weight or a doublet form can be observed in some cell types and/or following cell exposure to stressors. A synaptosome associated form of 40 kDa has recently been described.

## Notable Publications

Author	Pubmed ID	Journal	Application
Wenqian Wu	34679684	Antioxidants (Basel)	WB
Martin Kulessa	31545528	Neuropathol Appl Neurobiol	
Simona Cugusi	36121223	Mol Cell Biol	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10599-1-AP (BAG3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




IP result of anti-BAG3 (IP:10599-1-AP, 4ug; Detection:10599-1-AP 1:1000) with K-562 cells lysate 11000ug.




Immunofluorescent analysis of (4% PFA) fixed A549 cells using BAG3 antibody (10599-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using BAG3 antibody (10599-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 10599-1-AP (BAG3 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 10599-1-AP (BAG3 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).