

For Research Use Only

CSNK2A2 Polyclonal antibody

Catalog Number: 10606-1-AP **3 Publications**



Basic Information

Catalog Number:

10606-1-AP

Size:

400 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0877

GenBank Accession Number:

BC008812

GeneID (NCBI):

1459

UNIPROT ID:

P19784

Full Name:

casein kinase 2, alpha prime polypeptide

Calculated MW:

41 kDa

Observed MW:

30-35 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IF 1:20-1:200

Applications

Tested Applications:

IF/ICC, IP, WB, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human

Cited Species:

mouse

Positive Controls:

WB: HepG2 cells, HEK-293 cells, HT-1080 cells, Jurkat cells

IP: Jurkat cells,

IF: HepG2 cells,

Background Information

CSNK2A2 (Casein kinase II subunit alpha') is also named as CK2A3, CK2A2, CSNK2A1, CK2 and belongs to the protein kinase superfamily. It can function as a component of the p53/TP53-dependent spindle assembly checkpoint (SAC) that maintains cyclin-B-CDK1 activity and G2 arrest in response to spindle damage during mitosis. It also can protect against ionizing radiation-induced apoptosis partly by phosphorylating and activating SIRT1. This protein has molecular weights between 35 kDa and 44 kDa (PMID:8622692).

Notable Publications

Author	Pubmed ID	Journal	Application
Dahyun Yu	35659303	Acta Neuropathol Commun	WB, IHC
Yan Zhang	31987502	Biochem Biophys Res Commun	WB
Rachel H Mansky	36867535	Cell Rep	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 10606-1-AP (CSNK2A2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-CSNK2A2 (IP:10606-1-AP, 4ug; Detection:10606-1-AP 1:1000) with Jurkat cells lysate 2800 ug.



Immunofluorescent analysis of HepG2 cells using 10606-1-AP (CSNK2A2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).