For Research Use Only

UFD1L Polyclonal antibody

Catalog Number: 10615-1-AP

Featured Product

20 Publications

BC005087

Full Name:

Calculated MW: 35 kDa

(veast)

GenBank Accession Number:

ubiquitin fusion degradation 1 like



Basic Information

Applications

Catalog Number: 10615-1-AP

Isotype:

GeneID (NCBI): Size: 750 ug/ml 7353 Source: **UNIPROT ID:** Rabbit Q92890

IgG

Immunogen Catalog Number:

AG0954

Observed MW: 35 kDa

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Cited Applications: WB, IHC, IF, IP Species Specificity: human, mouse **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:50-1:500

Positive Controls:

WB: K-562 cells, mouse thymus tissue, Raji cells,

HepG2 cells, human testis tissue

IP: Raji cells,

IHC: human stomach tissue, IF/ICC: HepG2 cells, HeLa cells

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Michelle Harreman Lehner	36288698	Cell Rep	IF,
Xiaonan Wang	33037085	Mol Cancer Res	IHC
Yalcin Erzurumlu	36445513	Mol Biol Rep	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




K-562 cells were subjected to SDS PAGE followed by western blot with 10615-1-AP (UFD1L antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




IP result of anti-UFD1L (IP:10615-1-AP, 4ug; Detection:10615-1-AP 1:1000) with Raji cells lysate 2000ug.




Immunofluorescent analysis of HeLa cells using 10615-1-AP (UFD1L antibody) at dilution of 1:25 and Alexa Fluor 594-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using UFD11 antibody (10615-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 HEK-293 cells were intracellularly stained with 0.4 ug Anti-Human UFD1L (10615-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 10615-1-AP (UFD1L antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).