## For Research Use Only

# IL-18 Polyclonal antibody

Catalog Number: 10663-1-AP

Featured Product

328 Publications

GenBank Accession Number:



**Basic Information** 

Catalog Number: 10663-1-AP

Concentration: 750 ug/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG1063

Calculated MW: 193 aa, 22 kDa Observed MW:

22 kDa

BC007461

3606

II-18

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

Q14116

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:16000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF/ICC 1:50-1:500

**Applications** 

**Tested Applications:** 

WB, IF/ICC, FC (Intra), IP, ELISA

Cited Applications: WB, IHC, IF, IP Species Specificity: human, mouse Cited Species: human, mouse, rat **Positive Controls:** 

WB: A431 cells, HeLa cells, NIH/3T3 cells, HaCaT cell,

HaCaT cells, DU 145 cells

IP : HeLa cells,
IF/ICC : HeLa cells,

# **Background Information**

IL18, is a proinflammatory cytokine involved in the development of Th1 cells and in immune response. It can stimulate the NK cells and certain T cells to release IFN gamma which plays an important role in activating the macrophages or other cells. IL18 has been demonstrated to have the potential to enhance Fas ligand-mediated cytotoxicity, which is increased in PE and regulates placental apoptosis. IL18 is synthesized as a 22 kDa precursor and then cleaved into a biologically activate 18 kDa form. This antibody 10663-1-AP is raised against the full length IL18 precursor, and it recognizes both the pre-IL18 and mature IL18 theoretically.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
JiangFan Yu	36248980	Front Oncol	WB
Zhifang Li	34559953	CNS Neurosci Ther	WB
Deepavali Chakravarti	32958778	Nat Commun	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 10663-1-AP (IL-18 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using IL-18 antibody (10663-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



WB result of IL-18 antibody (10663-1-AP; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-IL-18 transfected HeLa cells.



1X10^6 HeLa cells were intracellularly stained with 0.2 ug Anti-Human IL-18 (10663-1-AP) and CoraLite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgC(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



IP result of anti-IL-18 (IP:10663-1-AP, 4ug; Detection:10663-1-AP 1:5000) with HeLa cells lysate 1360 ug.