## For Research Use Only

# ING5 Polyclonal antibody

Catalog Number: 10665-1-AP

**Featured Product** 

26 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 10665-1-AP BC005370
Size: GeneID (NCBI): 500 µg/ml 84289
Source: UNIPROT ID: Rabbit Q8WYH8

Isotype: Full Name:
IgG inhibitor of growth family, member 5

Immunogen Catalog Number: Calculated MW:

AG1059 28 kDa

Observed MW: 28-32 kDa

**Applications** 

Tested Applications: IHC, IP, WB, ELISA Cited Applications: IF, IHC, WB Species Specificity: human, mouse, rat

Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

WB: Jurkat cells, PC-3 cells, mouse kidney tissue, HEK-293 cells

**Purification Method:** 

WB 1:500-1:1000

protein lysate

IHC 1:20-1:200

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

IP: HEK-293 cells,

IHC: human colon cancer tissue, human ovary tumor

tissue

# **Background Information**

ING5, also named as p28ING5, belongs to the ING family. It is a component of the HBO1 complex which has a histone H4-specific acetyltransferase activity, a reduced activity toward histone H3 and is responsible for the bulk of histone H4 acetylation in vivo. ING5 is a component of the MOZ/MORF complex which has a histone H3 acetyltransferase activity. Through chromatin acetylation it may regulate DNA replication and may function as a transcriptional coactivator. (PMID:12750254) Altered ING5 expression and its shift into the cytoplasm may have an impact on the malignant transformation of colorectal epithelial cells and should be considered as a biomarker for colorectal carcinogenesis. (PMID:21193223)

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yahan Wu	32973339	Cell Res	WB
Kumi Matsuura	32918898	Exp Cell Res	WB
Neslisah Barlak	31475765	Cell Biol Int	WB

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.cor

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


# **Selected Validation Data**




WB result of ING5 antibody (10665-1-AP; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ING5 transfected Jurkat cells.



Jurkat cells were subjected to SDS PAGE followed by western blot with 10665-1-AP (ING5 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 burs.



Immunohistochemical analysis of paraffinembedded human colon cancer using 10665-1-AP (ING5 antibody) at dilution of 1:100 (under 25x lens)



IP result of anti-ING5 (IP:10665-1-AP, 4ug; Detection:10665-1-AP 1:700) with HEK-293 cells lysate 2560ug.