

For Research Use Only						
VAMP3/Cellubrevin Polyclonal proteintech [®]						
antibody	, , , , , , , , , , , , , , , , , , ,			Antibodies ELISA kits Proteins WWW.ptglab.com		
Catalog Number:10702-1-A	Featured Product	16 Publicatior	ns			
Basic Information	Catalog Number: 10702-1-AP	GenBank Accession Number: BC005941 GeneID (NCBI): 9341 UNIPROT ID: Q15836 Full Name: vesicle-associated membrane protei 3 (cellubrevin)		Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:20-1:200 rotein		
	Size: 700 µg/ml					
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG1156					
					Calculated MW: 11 kDa	
		Observed MW: 11-17 kDa				
		Applications	Tested Applications:		Positive Controls:	
IHC, IP, WB,ELISA Cited Applications: WB, IHC, Blocking, IF	WB : HEK-293 cells, MCF-7 cells, mouse brain tissue					
	IP : HEK-293 cells,					
Species Specificity: IHC : human heart tissue, human kidney tissue human, mouse, rat Cited Species: human, mouse, rat Image: Species in the species i						
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	VAMP3, also named as cellubrevin or synaptobrevin 3, is a member of the vesicle-associated membrane protein (VAMP)/synaptobrevin family. VAMP3 is a v-SNARE (soluble NSF-attachment protein receptor). Characterized by a common sequence called the SNARE motif, SNARE proteins are involved in membrane fusion and vesicular transport (PMID: 11252968). VAMP3 resides in recycling endosomes and endosome-derived transport vesicles, involved in regulating membrane traffic. It is implicated in recycling of transferrin receptors, secretion of alpha-granules in platelets, and membrane trafficking during cell migration.					
Notable Publications	Author	Pubmed ID Jo	ournal	Application		
	Yonglu Che	31712554 N	at Commun	WB,IF		
	Juan-Juan Zhu	33224946 Fi	ront Cell Dev Biol	IF		
	Xiaoli Wang	29785746 Pi	roteomics	WB		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	1 50% glycerol pH 7.3.				

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data






WB result of VAMP3/Cellubrevin antibody (10702-
1-AP; 1:2000; incubated at room temperature for 1.5
hours) with sh-Control and sh-VAMP3/Cellubrevin
transfected HEK-293 cells.HEK-293 cells were subjected to SDS PAGE
followed by western blot with 10702-1-AP
(VAMP3/Cellubrevin antibody) at dilution of
1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-VAMP3/Cellubrevin (IP:10702-1-AP, 4ug; Detection:10702-1-AP 1:1000) with HEK-293 cells lysate 3000ug.



Immunohistochemical analysis of paraffinembedded human heart tissue slide using 10702-1-AP (VAMP3/Cellubrevin antibody at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human heart tissue slide using 10702-1-AP (VAMP3/Cellubrevin antibody at dilution of 1:50 (under 40x lens).