For Research Use Only

CXCL11 Polyclonal antibody

Catalog Number: 10707-1-AP 2 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 10707-1-AP
 BC005292

 Size:
 GeneID (NCBI):

 700 ug/ml
 6373

 Source:
 UNIPROT ID:

 Rabbit
 014625

Full Name: chemokine (C-X-C motif) ligand 11

Immunogen Catalog Number: Calculated MW:

AG1087 10 kDa

Observed MW: 10 kDa

Applications

Tested Applications: WB, IHC, ELISA Cited Applications:

IHC

Isotype:

Species Specificity:

human Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: IFN gamma, LPS and Brefeldin A treated THP-1

Purification Method:

WB 1:500-1:1000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

cells,

IHC: human colon cancer tissue,

Background Information

CXCL11, also known as IFN-inducible T-cell $\,^{\alpha}$ -chemoattractant (I-TAC), is a member of the ELR-negative CXC chemokine family. It is produced by various cells including leukocytes, fibroblasts, and endothelial cells upon stimulation with interferons (IFNs). CXCL11 signals through the chemokine receptors CXCR3 and CXCR7 . This chemokine is associated with multiple functions such as chemotactic migration, regulation of cell proliferation and self-renewal, increasing cell adhesion, and modulation of angiostatic effects.

Notable Publications

Author	Pubmed ID	Journal	Application
Haihang Nie	38834869	Cancer Immunol Immunother	IHC
Zihan Xi	38137348	Biomedicines	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 10707-1-AP (CXCL11 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IFN gamma, LPS and Brefeldin A treated THP-1 cells were subjected to SDS PAGE followed by western blot with 10707-1-AP (CXCL11 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.