For Research Use Only

CNOT8 Polyclonal antibody

Catalog Number: 10752-1-AP

Featured Product

7 Publications



Basic Information

Catalog Number: GenBank Accession Number: 10752-1-AP BC017366 GeneID (NCBI): Concentration: 700 ug/ml **UNIPROT ID:** Source: Rabbit Q9UFF9 Full Name: Isotype:

CCR4-NOT transcription complex,

subunit 8 Immunogen Catalog Number:

AG1201 Calculated MW:

34 kDa Observed MW: 34 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200 IF/ICC 1:200-1:800

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

Species Specificity: human, mouse **Cited Species:**

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, HeLa cells, mouse testis tissue

IP: mouse testis tissue,

IHC: human prostate cancer tissue, human kidney

tissue

IF/ICC: HeLa cells,

Background Information

CCR4-NOT transcription complex subunit 8(CNOT8), is a ubiquitous transcription factor, which required for a diverse set of processes. The CCR4-NOT complex is a global regulator of RNA polymerase II, and functions as general transcription regulation complex. It consisted part by CCR4, NOT1 to NOT5, and CAF1. This is a rabbit polyclonal antibody raised against the full-length chain of human CNOT8.

Notable Publications

Author	Pubmed ID	Journal	Application
Tian-Li Duan	31936572	Cells	WB
Jing Yang	39747175	Nat Commun	
Yusheng Liu	38956014	Nat Commun	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 10752-1-AP (CNOT8 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer using 10752-1-AP (CNOT8 antibody) at dilution of 1:100 (under 10x Jens).



IP result of anti-CNOT8 (IP:10752-1-AP, 4ug; Detection:10752-1-AP 1:600) with mouse testis tissue lysate 4000ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CNOT8 antibody (10752-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).