

For Research Use Only

SLC25A13 Polyclonal antibody

Catalog Number: 10789-1-AP



Basic Information

Catalog Number:

10789-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1242

GenBank Accession Number:

BC006566

GeneID (NCBI):

10165

UNIPROT ID:

Q9UJ50

Full Name:

solute carrier family 25, member 13 (citrin)

Calculated MW:

74 kDa

Observed MW:

60-63 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2400

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF 1:50-1:500

Applications

Tested Applications:

IF/ICC, IHC, IP, WB, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse kidney tissue, Raji cells, A431 cells, HEK-293 cells, mouse liver tissue

IP: mouse kidney tissue,

IHC: human colon cancer tissue,

IF: U2OS cells,

Background Information

SLC25A13, also known as citrin, has been identified as a mitochondrial aspartate-glutamate carrier protein and may be implicated in metabolic compartmentation. It consists of 675 amino acid residues and harbors four EF-hands and six mitochondrial transmembranous (TM) spanners. Mutations in SLC25A13 gene cause adult-onset type II citrullinemia (CTLN2) and neonatal intrahepatic cholestasis (NICCD). The complete deletion or cleaved SLA25A13 proteins have been reported in patients with SLC25A13 mutation.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




mouse kidney tissue were subjected to SDS PAGE followed by western blot with 10789-1-AP (SLC25A13 antibody) at dilution of 1:1200 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed U2OS cells using SLC25A13 antibody (10789-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 10789-1-AP (SLC25A13 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 10789-1-AP (SLC25A13 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-SLC25A13 (IP:10789-1-AP, 4 μ g; Detection:10789-1-AP 1:1000) with mouse kidney tissue lysate 4000 μ g.