## For Research Use Only

## ARF3 Polyclonal antibody

Catalog Number: 10800-1-AP 1 Publications



**Basic Information** 

Catalog Number: 10800-1-AP

Concentration: 500 ug/ml

Source: Rabbit Isotype:

gG ADP-ribosylation factor 3

Immunogen Catalog Number: Calculated MW:

AG1174 21 kDa

Observed MW: 18-20 kDa

BC017565

GeneID (NCBI):

**UNIPROT ID:** 

P61204 Full Name:

GenBank Accession Number:

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000 IHC 1:50-1:500 IF/ICC 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications:

WB, IHC

Species Specificity: human, mouse, rat Cited Species: mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: mouse brain tissue, HeLa cells, HepG2 cells, human brain tissue, mouse cerebellum tissue, Raji

cells, rat brain tissue

IF/ICC: HepG2 cells,

 $\label{eq:IHC:human} \textbf{IHC:} human \ kidney \ tissue, human \ pancreas \ tissue,$ 

human pancreas cancer tissue

Background Information

ADP-ribosylation factors (ARFs) are members of the ARF family of GTP-binding proteins of the Ras superfamily, with 20kda protein size. ARFs bind and regulate GTP/GDP cycle by alternating between the active GTP-bound and inactive GDP-bound conformations. ARF family proteins are essential and ubiquitous in eukaryotes. Six highly conserved members of the family have been identified in mammalian cells. They function in vesicular traffic and actin remodelling and other bioprocesses in cells. ARF3 is 95% homolouous to ARF1. Its function was not fully understood. (PMID: 7759471, PMID: 16042562). This antibody can bind ARFs for the close sequences.

**Notable Publications** 

 Author
 Pubmed ID
 Journal
 Application

 Naila Umer
 35950913
 Development
 WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**

 $70kd \rightarrow$   $50kd \rightarrow$   $40kd \rightarrow$   $30kd \rightarrow$   $20kd \rightarrow$   $15kd \rightarrow$   $10kd \rightarrow$   $5kd \rightarrow$ 


mouse brain tissue were subjected to SDS PAGE followed by western blot with 10800-1-AP (ARF3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 10800-1-AP (ARF3 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 10800-1-AP (ARF3 Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using ARF3 antibody (10800-1-AP) at dilution of 1:200 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4).