For Research Use Only

CD69 Polyclonal antibody

Catalog Number: 10803-1-AP 12 Publications



Basic Information

Catalog Number: 10803-1-AP

Concentration: 750 ug/ml

Source: Rabbit Isotype:

Immunogen Catalog Number: AG1094

23 kDa

Observed MW: 28 kDa, 32 kDa

CD69 molecule Calculated MW:

GenBank Accession Number:

BC007037

GeneID (NCBI):

UNIPROT ID:

Q07108 Full Name: **Purification Method:**

Antigen affinity purification

Recommended Dilutions: WB: 1:500-1:3000 IHC: 1:1000-1:4000 IF-P: 1:50-1:500

Applications

Tested Applications: WB, IHC, IF-P, ELISA Cited Applications: WB, IHC, IF

Species Specificity:

human **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human peripheral blood platelets,

IHC: human tonsillitis tissue, IF-P: human tonsillitis tissue,

Background Information

CD69, also known as activation inducer molecule (AIM) or early activation antigen (EA1), is a cell surface glycoprotein expressed shortly after activation on T lymphocytes and other cells of haematopoietic origin, including natural killer cells. It is a disulfide-linked homodimer composed of two chains belonging to the type II integral protein with a C-type lectin domain. CD69 is involved in lymphocyte proliferation and functions as a signaltransmitting receptor in lymphocytes, natural killer cells, and platelets.

Notable Publications

Author	Pubmed ID	Journal	Application
Shijia Rao	31661988	Expert Rev Clin Immunol	IF
Huiwen Zheng	35594870	Cell Rep	IF
Chen Zhu	31428519	Oncoimmunology	IHC

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 10803-1-AP (CD69 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 10803-1-AP (CD69 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 10803-1-AP (CD69 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 10803-1-AP (CD69 antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 10803-1-AP (CD69 antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD69 antibody (10803-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).