For Research Use Only

LIX/CXCL5 Polyclonal antibody

Catalog Number: 10809-1-AP 1 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 10809-1-AP
 BC008376

 Size:
 GeneID (NCBI):

 1000 ug/ml
 6374

 Source:
 UNIPROT ID:

Source: UNIPROTIE
Rabbit P42830
Isotype: Full Name:

IgG chemokine (C-X-C motif) ligand 5

Immunogen Catalog Number: Calculated MW:

AG1079 12 kDa

Observed MW: 8 kDa

Antigen affinity purification Recommended Dilutions: WB 1:200-1:1000 IHC 1:200-1:800

Purification Method:

Applications

Tested Applications: WB, IHC, ELISA Cited Applications:

cited Applications

WB

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: PMA treated A549 cells,

IHC: human intrahepatic cholangiocarcinoma tissue,

Background Information

Chemokines are a class of pro-inflammatory cytokines that can recruit and activate chemotactic cells. CXCL5 is a member of the chemokine family binding CXCR2, a G-protein coupled receptor. CXCL5 is a chemokine that promotes tumor formation by triggering the migration of immune cells to tumors and promotes immuno-suppressive characteristics of the tumor microenvironment. CXCL5 can also promote tumor cell metastasis and recruit vascular endothelial cells for angiogenesis.

Notable Publications

Author	Pubmed ID	Journal	Application
Peng Huang	39428975	Stem Cells	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Untreated, PMA treated A549 cells were subjected to SDS PAGE followed by western blot with 10809-1-AP (CXCL5 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human intrahepatic cholangiocarcinoma tissue slide using 10809-1-AP (CXCL5 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).