

For Research Use Only

ERCC2 Polyclonal antibody

Catalog Number: 10818-1-AP

14 Publications



Basic Information

Catalog Number:

10818-1-AP

Size:

300 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1038

GenBank Accession Number:

BC008346

GeneID (NCBI):

2068

UNIPROT ID:

P18074

Full Name:

excision repair cross-complementing
rodent repair deficiency,
complementation group 2

Calculated MW:

87 kDa

Observed MW:

80 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total
protein lysate

IHC 1:20-1:200

Applications

Tested Applications:

WB, IP, IHC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human

Cited Species:

human, mouse, rat

Positive Controls:

WB : HeLa cells, K-562 cells, HEK-293 cells

IP : HEK-293 cells,

IHC : human lymphoma tissue, human cervical cancer
tissue

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Background Information

Excision-repair complementing defective in Chinese hamster 2(ERCC2), belongs to the nucleotide excision repair pathway, is a component of the TFIIH transcription complex and required for the association of the CDK-activating kinase(CAK) subcomplex with TFIIH. Since TFIIH is involved in both DNA repair and cell cycle progression, ERCC2 is implicated to play an integral role in the nucleotide excision repair pathway. Also it involves in the regulation of vitamin-D receptor activity. As a part of the mitotic spindle-associated MMXD complex, ERCC2 has a role in chromosome segregation.

Notable Publications

Author	Pubmed ID	Journal	Application
Shun Tezuka	30416859	Am J Cancer Res	IHC
Lai Jiun-I JI	19432884	Cancer Sci	IHC
Aihua Zhang	27685703	Biomarkers	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 10818-1-AP (ERCC2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lymphoma using 10818-1-AP (ERCC2 antibody) at dilution of 1:100 (under 10x lens).



IP result of anti-ERCC2 (IP:10818-1-AP, 4ug; Detection:10818-1-AP 1:1000) with HEK-293 cells lysate 1575 ug.