

For Research Use Only

# c-MYC Polyclonal antibody

Catalog Number: 10828-1-AP

Featured Product

706 Publications



## Basic Information

### Catalog Number:

10828-1-AP

### Concentration:

800 ug/ml

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG1263

### GenBank Accession Number:

BC000141

### GeneID (NCBI):

4609

### UNIPROT ID:

P01106

### Full Name:

v-myc myelocytomatosis viral oncogene homolog (avian)

### Calculated MW:

49 kDa

### Observed MW:

62-65 kDa, 50 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:2000-1:12000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:500-1:2000

IF/ICC 1:200-1:800

## Applications

### Tested Applications:

WB, IF/ICC, FC (Intra), IP, ELISA

### Cited Applications:

WB, IF, IP, CoIP, ChIP, RIP, EMSA

### Species Specificity:

human

### Cited Species:

human, mouse, rat, pig, chicken, sheep, p. lilacinum

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB** : HT-29 cells, Jurkat cells, HeLa cells, A549 cells, HL-60 cells, human placenta tissue, Raji cells, SH-SY5Y cells, HepG2 cells, BxPC-3 cells, SW 1990 cells, Jurkat cells and Raji cells.

**IP** : MCF-7 cells,

**IHC** : human ovary cancer tissue,

**IF/ICC** : Saos-2 cells, HEK-293 cells, U2OS cells

## Background Information

MYC contains one basic helix-loop-helix (bHLH) domain. This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. It seems to activate the transcription of growth-related genes. It binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. This antibody is a rabbit polyclonal antibody raised against recombinant protein of human MYC. The 50kDa band recognized by antibody is the native form of MYC, while the other bands, between 60-70kDa, are the phosphorylated form of MYC [PMID: 12189186]

## Notable Publications

Author	Pubmed ID	Journal	Application
Ruosi Yao	33061303	Drug Des Devel Ther	WB
Zhen-Yu Shi	29039446	Int J Mol Med	WB
Ke-Xin Wang	34649212	Phytomedicine	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


Selected Validation Data




WB result of c-MYC antibody (10828-1-AP; 1:12000; incubated at room temperature for 1.5 hours) with sh-Control and sh-c-MYC transfected A549 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 10828-1-AP (c-MYC antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




IP result of anti-c-MYC (IP:10828-1-AP, 4ug; Detection:10828-1-AP 1:1000) with MCF-7 cells lysate 1660 ug.




Immunohistochemical analysis of paraffin-embedded human ovary cancer tissue slide using 10828-1-AP (c-MYC antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed Saos-2 cells using c-MYC antibody (10828-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



1x10<sup>6</sup> HeLa cells were intracellularly stained with 0.4 ug Anti-Human c-MYC (10828-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



1x10<sup>6</sup> NCCIT cells were intracellularly stained with 0.25 ug c-MYC Polyclonal antibody (10828-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



1x10<sup>6</sup> HeLa cells were intracellularly stained with 0.25 ug c-MYC Polyclonal antibody (10828-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).