

For Research Use Only

# NOL3 Polyclonal antibody

Catalog Number: 10846-2-AP

7 Publications



## Basic Information

Catalog Number:

10846-2-AP

Concentration:

1100 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1306

GenBank Accession Number:

BC012798

GeneID (NCBI):

8996

UNIPROT ID:

O60936

Full Name:

nucleolar protein 3 (apoptosis repressor with CARD domain)

Calculated MW:

24 aa, 3 kDa

Observed MW:

30-33 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:50-1:500

IF/ICC 1:200-1:800

## Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IHC, IF, IP

Species Specificity:

human, mouse

Cited Species:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

WB : mouse skeletal muscle tissue, HeLa cells

IHC : human breast cancer tissue,

IF/ICC : HepG2 cells,

## Background Information

### Notable Publications

Author	Pubmed ID	Journal	Application
R Zajackowska	30267794	Pharmacol Biochem Behav	WB
X Li	25341033	Cell Death Dis	WB
Zhen Zhang	33193701	Front Genet	IHC

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10846-2-AP (NOL3 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 10846-2-AP (NOL3 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using NOL3 antibody (10846-2-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).