

For Research Use Only

# MECP2 Polyclonal antibody

Catalog Number: 10861-1-AP

Featured Product

22 Publications



## Basic Information

Catalog Number:

10861-1-AP

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1309

GenBank Accession Number:

BC011612

GeneID (NCBI):

4204

UNIPROT ID:

P51608

Full Name:

methyl CpG binding protein 2 (Rett syndrome)

Calculated MW:

52 kDa, 53 kDa

Observed MW:

75 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:1000-1:4000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:50-1:500

IF-P: 1:50-1:500

FC (Intra): 0.25 ug per 10<sup>6</sup> cells in a 100 µl suspension

## Applications

Tested Applications:

WB, IHC, IF-P, IP, ELISA

Cited Applications:

WB, IHC, IF, ChIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

WB : MCF-7 cells, mouse lung tissue, mouse brain tissue, rat lung tissue

IP : HEK-293T cells, MCF-7 cells

IHC : human gliomas tissue, human breast cancer tissue, rat brain tissue

IF-P : mouse cerebellum tissue,

FC (Intra) : NIH/3T3 cells,

## Background Information

METHYL-CpG-BINDING PROTEIN 2 (MECP2), is a chromatin-associated protein that can both activate and repress transcription. Mecp2 in the adult mouse is high in the brain, lung, and spleen, lower in heart and kidney. MECP2 takes part in the control of neuronal activity-dependent gene regulation, and this process may underlie the pathology of Rett syndrome, a severe developmental disorder with autistic phenotypes. As compared to wild-type monkeys, MECP2 transgenic monkeys exhibited a higher frequency of repetitive circular locomotion and increased stress responses, as measured by the threat-related anxiety and defensive test. Chromosomal protein that binds to methylated DNA. It can bind specifically to a single methyl-CpG pair. It is not influenced by sequences flanking the methyl-CpGs. Mediates transcriptional repression through interaction with histone deacetylase and the corepressor SIN3A. Two isoforms of MECP2 exist due to alternative splicing events. This antibody reacts with the MECP2 and phosphorylated MECP2 proteins. The calculated molecular weight of MECP2 is a 52 kDa, but the post-modified MECP2 protein is about 75-80 kDa (PMID: 12160743).

## Notable Publications

Author	Pubmed ID	Journal	Application
Gamze Ayaz	34475492	Sci Rep	ChIP
Wang Ying Y	21925646	Brain Res	WB, IHC
Jordan M Buck	32138755	Epigenetics Chromatin	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




IP result of anti-MECP2 (IP:10861-1-AP, 4ug; Detection:10861-1-AP 1:500) with HEK-293T cells lysate 1640 ug.




Various lysates were subjected to SDS PAGE followed by western blot with 10861-1-AP (MECP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 10861-1-AP (MECP2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 10861-1-AP (MECP2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 10861-1-AP (MECP2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 10861-1-AP (MECP2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse cerebellum tissue using MECP2 antibody (10861-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse cerebellum tissue using MECP2 antibody (10861-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1x10<sup>6</sup> NIH/3T3 cells were intracellularly stained with 0.25 ug MECP2 Polyclonal antibody (10861-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug rabbit IgG isotype control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).