For Research Use Only

Cyclin G Polyclonal antibody

Catalog Number: 10897-1-AP

5 Publications



Basic Information

Catalog Number: 10897-1-AP Size:

260 µg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG1157

Observed MW: 34 kDa

BC007093

GeneID (NCBI):

UNIPROT ID:

P51959 Full Name:

cyclin G1 Calculated MW:

34 kDa

GenBank Accession Number:

Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500

Applications

Tested Applications: IHC, WB, ELISA Cited Applications:

WB

Species Specificity: human, mouse, rat **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: MDA-MB-453s cells, mouse ovary tissue, HeLa

cells, mouse liver tissue

IHC: human breast cancer tissue,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Fanpeng Zhao	25342750	J Biol Chem	WB
Yuzhong Chen	36400182	Exp Cell Res	WB
Chen-Song Huang	29458145	Cancer Lett	WB

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



MDA-MB-453s cells were subjected to SDS PAGE followed by western blot with 10897-1-AP (Cyclin G antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10897-1-AP (Cyclin G antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10897-1-AP (Cyclin G antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).