

For Research Use Only

AMPK Alpha Polyclonal antibody

Catalog Number: 10929-2-AP

Featured Product

261 Publications



Basic Information

Catalog Number:

10929-2-AP

Concentration:

500 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1367

GenBank Accession Number:

BC012622

GeneID (NCBI):

5562

UNIPROT ID:

Q13131

Full Name:

protein kinase, AMP-activated, alpha
1 catalytic subunit

Calculated MW:

63 kDa

Observed MW:

64 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:10000

IP 0.5-4.0 ug for 1.0-3.0 mg of total
protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat, pig, rabbit, zebrafish, goat, fish,
carp

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : HeLa cells, Jurkat cells, NIH/3T3 cells

IP : HeLa cells,

IHC : human prostate cancer tissue,

IF/ICC : HepG2 cells,

Background Information

The mammalian 5-prime-AMP-activated protein kinase (AMPK) appears to play a role in protecting cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. PRKAA1 is also named as AMPK1, ACACA kinase, HMGCR kinase. It is a mammalian homologue of sucrose non-fermenting protein kinase (SNF-1), which belongs to a serine/threonine protein kinase family. It has 2 isoforms with molecular mass of 63-66 kDa produced by alternative splicing. 10929-2-AP can recognize both AMPK Alpha 1 and AMPK Alpha 2.

Notable Publications

Author	Pubmed ID	Journal	Application
Xiao-Fei Guo	29113747	Prostaglandins Leukot Essent Fatty Acids	WB
Xiaochen Bai	34543233	Aging (Albany NY)	WB
Shijie Cao	30268048	Environ Toxicol Pharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10929-2-AP (AMPK Alpha 1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 10929-2-AP (AMPK Alpha 1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 10929-2-AP (AMPK Alpha 1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-AMPK Alpha (IP:10929-2-AP, 4ug; Detection:10929-2-AP 1:1000) with HeLa cells lysate 1600 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using AMPK Alpha antibody (10929-2-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).