For Research Use Only

Galectin 8 Polyclonal antibody

Catalog Number:10955-1-AP





Basic Information	Catalog Number: 10955-1-AP	GenBank Accession Number BC015818	: Purification Method: Antigen affinity purification	
	Size: 300 ug/ml Source: Rabbit	GeneID (NCBI): 3964 UNIPROT ID: 000214	Recommended Dilutions: WB 1:500-1:1000 IHC 1:200-1:800 IF/ICC 1:200-1:800	
				Isotype: Full Name: IgG lectin, galactoside-binding, soluble, 8
	Immunogen Catalog Number: AG1387	Calculated MW: 34 kDa		
		Observed MW: 36 kDa, 40 kDa		
	Applications			tive Controls:
WB : HEK-293 cells, LNCaP cells				
human			IHC : human intrahepatic cholangiocarcinoma tissue,	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		C : HEK-293 cells,		
Background Information	Galectin 8 is a member of the galectin family. Galectins are beta-galactoside-binding animal lectins with conserved carbohydrate recognition domains. The galectins have been implicated in many essential functions including development, differentiation, cell-cell adhesion, cell-matrix interaction, growth regulation, apoptosis, and RNA splicing. This gene is widely expressed in tumoral tissues and seems to be involved in integrin-like cell interactions.			
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	and 50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded human intrahepatic cholangiocarcinoma tissue slide using 10955-1-AP (Galectin 8 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using Galectin 8 antibody (10955-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).

Various lysates were subjected to SDS PAGE followed by western blot with 10955-1-AP (Galectin 8 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.