For Research Use Only

T: 4006900926

E: Proteintech-CN@ptglab.com

CSN2 Polyclonal antibody Catalog Number:10969-2-AP Featured Product

Featured Product

10 Publications




Group brand and is not available to purchase from any

other manufacturer.


Basic Information	Catalog Number: 10969-2-AP	GenBank Accession N BC012629	lumber:	Purification Method: Antigen affinity purification	
	Concentration:	GeneID (NCBI):		Recommended Dilutions:	
	600 ug/ml	9318		WB 1:500-1:1000	
	Source:	UNIPROT ID:		IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	Rabbit Isotype:	P61201 Full Name:		IHC 1:50-1:500	
	IgG Immunogen Catalog Number:	IF/ICC 1:50-1:500 COP9 constitutive photomorphogenic homolog subunit 2 (Arabidopsis) Calculated MW: 52 kDa			
	AG1415				
		Observed MW: 52 kDa			
Applications	Tested Applications:			ols:	
	WB, IHC, IF/ICC, IP, ELISA		WB : HeLa cells, HEK-293T cells, HEK-293 cells, mou		
	Cited Applications: WB, IHC, IF			in tissue, NIH/3T3 cells, mouse kidney tissue	
	Species Specificity:		IP : NIH/3T3 cells,		
	human, mouse		IHC : human ovary tumor tissue, IF/ICC : HeLa cells,		
	Cited Species: human, mouse				
	TE buffer pH 9.0; (*) Altern	atively, antigen			
	TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0				
Background Informatio	 retrieval may be performed buffer pH 6.0 COPS2 is an essential component and developmental processes. Th pathway by mediating the dened decrease the Ubl ligase activity of phosphorylation of p53/TP53, crju with CK2 and PKD kinases. CSN-d the Ubl system, respectively. (reference) 	d with citrate t of the COP9 signalosome te CSN complex is an esse dylation of the cullin subu of SCF-type complexes suc un/JUN, IkappaBalpha/NF tependent phosphorylation er to UniProt) Catalog#10	ntial regulator of inits of SCF-type h as SCF, CSA or KBIA, ITPK1 and I n of TP53 and JUN 969-2-AP is a rab		
	retrieval may be performed buffer pH 6.0 COPS2 is an essential component and developmental processes. Th pathway by mediating the dened decrease the Ubl ligase activity of phosphorylation of p53/TP53, c-ju with CK2 and PKD kinases. CSN-d the Ubl system, respectively. (refit the full-length of human COPS2.	d with citrate t of the COP9 signalosome te CSN complex is an esse dylation of the cullin subu of SCF-type complexes suc un/JUN, IkappaBalpha/NF tependent phosphorylation er to UniProt) Catalog#10	ntial regulator of nits of SCF-type h as SCF, CSA or KBIA, ITPK1 and I n of TP53 and JUN 969-2-AP is a rab 12 kDa, and this a	the ubiquitin (Ubl) conjugation E3 ligase complexes, leading to DDB2. The complex is also involved RF8/ICSBP, possibly via its associati I promotes and protects degradation ubit polyclonal antibody raised again	
	retrieval may be performed buffer pH 6.0 ON COPS2 is an essential component and developmental processes. Th pathway by mediating the dened decrease the Ubl ligase activity o phosphorylation of p53/TP53, c-ju with CK2 and PKD kinases. CSN-d the Ubl system, respectively. (refe the full-length of human COPS2." protein.	d with citrate to f the COP9 signalosome the CSN complex is an essed dylation of the cullin subu of SCF-type complexes suc un/JUN, IkappaBalpha/NF bependent phosphorylation er to UniProt) Catalog#10 The MW of this protein is 5 Pubmed ID Journ	ntial regulator of nits of SCF-type h as SCF, CSA or KBIA, ITPK1 and I n of TP53 and JUN 969-2-AP is a rab 12 kDa, and this a	the ubiquitin (Ubl) conjugation E3 ligase complexes, leading to DDB2. The complex is also involved RF8/ICSBP, possibly via its associati I promotes and protects degradation ubit polyclonal antibody raised again ntibody specially recognises the 52 l Application	
	retrieval may be performed buffer pH 6.0 COPS2 is an essential component and developmental processes. Th pathway by mediating the dened decrease the Ubl ligase activity of phosphorylation of p53/TP53, c-ju with CK2 and PKD kinases. CSN-d the Ubl system, respectively. (refit the full-length of human COPS2. protein.	d with citrate t of the COP9 signalosome he CSN complex is an esse dylation of the cullin subu of SCF-type complexes suc un/JUN, IkappaBalpha/NF ependent phosphorylation er to UniProt) Catalog#10 The MW of this protein is 5 Pubmed ID Journ 25349427 Proc	ntial regulator of nits of SCF-type h as SCF, CSA or KBIA, ITPK1 and I of TP53 and JUN 969-2-AP is a rab i2 kDa, and this a	the ubiquitin (Ubl) conjugation E3 ligase complexes, leading to DDB2. The complex is also involved RF8/ICSBP, possibly via its associati I promotes and protects degradation bit polyclonal antibody raised again ntibody specially recognises the 52 l Application A WB	
Background Informatio	retrieval may be performed buffer pH 6.0 COPS2 is an essential component and developmental processes. Th pathway by mediating the dened decrease the Ubl ligase activity o phosphorylation of p53/TP53, c-ju with CK2 and PKD kinases. CSN-d the Ubl system, respectively. (ref the full-length of human COPS2. protein. Author Feng Rao	d with citrate t of the COP9 signalosome te CSN complex is an esse dylation of the cullin subu of SCF-type complexes suc un/JUN, IkappaBalpha/NF ependent phosphorylation er to UniProt) Catalog#10 The MW of this protein is 5 Pubmed ID Journ 25349427 Proc 26976604 Proc	ntial regulator of inits of SCF-type h as SCF, CSA or KBIA, ITPK1 and I of TP53 and JUN 969-2-AP is a rab i2 kDa, and this a nal Natl Acad Sci U S	the ubiquitin (Ubl) conjugation E3 ligase complexes, leading to DDB2. The complex is also involved RF8/ICSBP, possibly via its associati I promotes and protects degradation ubit polyclonal antibody raised again ntibody specially recognises the 52 Application A WB	
	retrieval may be performed buffer pH 6.0 COPS2 is an essential component and developmental processes. Th pathway by mediating the dened decrease the Ubl ligase activity of phosphorylation of p53/TP53, c-jt with CK2 and PKD kinases. CSN-d the Ubl system, respectively. (refit the full-length of human COPS2.) protein. Author Feng Rao Paul C Scherer	d with citrate t of the COP9 signalosome he CSN complex is an esse dylation of the cullin subu of SCF-type complexes suc un/JUN, IkappaBalpha/NF ependent phosphorylation er to UniProt) Catalog#10 The MW of this protein is 5 Pubmed ID Journ 25349427 Proc 26976604 Proc 32047038 Proc	ntial regulator of nits of SCF-type h as SCF, CSA or KBIA, ITPK1 and I of TP53 and JUN 969-2-AP is a rab i2 kDa, and this a nal Natl Acad Sci U S	the ubiquitin (Ubl) conjugation E3 ligase complexes, leading to DDB2. The complex is also involved RF8/ICSBP, possibly via its associati I promotes and protects degradation ubit polyclonal antibody raised again ntibody specially recognises the 52 l Application A WB A WB	

W: ptgcn.com


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10969-2-AP (CSN2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 10969-2-AP (CSN2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 10969-2-AP (CSN2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




IP result of anti-CSN2 (IP:10969-2-AP, 3ug; Detection:10969-2-AP 1:500) with NIH/3T3 cells lysate 2500ug.



HeLa cells were subjected to SDS PAGE followed by western blot with 10969-2-AP (CSN2 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



WB result of CSN2 antibody (10969-2-AP; 1:1200; incubated at room temperature for 1.5 hours) with sh-Control and sh-CSN2 transfected HEK-293 cells.



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using CSN2 antibody (10969-2-AP) at dilution of 1:200 and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L).