

For Research Use Only

EIF2B2 Polyclonal antibody

Catalog Number: 11034-1-AP

Featured Product

9 Publications



Basic Information

Catalog Number:

11034-1-AP

Size:

450 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1502

GenBank Accession Number:

BC011750

GeneID (NCBI):

8892

UNIPROT ID:

P49770

Full Name:

eukaryotic translation initiation factor 2B, subunit 2 beta, 39kDa

Calculated MW:

39 kDa

Observed MW:

39 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:14000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications:

WB, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, mouse liver tissue, HEK-293 cells, K-562 cells

IP: HEK-293 cells,

IHC: rat brain tissue, mouse brain tissue

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Michael Schoof	33688831	Elife	WB
Noel C Wortham	26614765	Biochem J	WB
Yu Lin	31215738	Mol. Reprod Dev	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




A431 cells were subjected to SDS PAGE followed by western blot with 11034-1-AP (EIF2B2 antibody) at dilution of 1:7000 incubated at room temperature for 1.5 hours.




IP result of anti-EIF2B2 (IP:11034-1-AP, 4ug; Detection:11034-1-AP 1:500) with HEK-293 cells lysate 2000 ug.




Various lysates were subjected to SDS PAGE followed by western blot with 11034-1-AP (EIF2B2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 11034-1-AP (EIF2B2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 11034-1-AP (EIF2B2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 11034-1-AP (EIF2B2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 11034-1-AP (EIF2B2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).