#### For Research Use Only

# **UXT Polyclonal antibody**

Catalog Number: 11047-1-AP

1 Publications



**Purification Method:** 

WB 1:200-1:1000

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

**Basic Information** 

Catalog Number: 11047-1-AP Size:

 $600 \mu g/ml$ Source: Rabbit Isotype:

Immunogen Catalog Number:

AG1513

18 kDa, 45 kDa

20 kDa

GenBank Accession Number:

BC008890 GeneID (NCBI): 8409

**UNIPROT ID:** Q9UBK9 Full Name:

Calculated MW:

ubiquitously-expressed transcript

Observed MW:

**Applications** 

**Tested Applications:** 

IP, WB, ELISA Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species: mouse

Positive Controls:

WB: HeLa cells, mouse brain tissue

IP: mouse brain tissue.

## **Background Information**

Ubiquitously expressed transcript (UXT) is a gene transcription regulator. It can regulate androgen receptor(AR) transcription by concerting with the coreperssor UR1. It is proposed a potential component of mitochondrialassociated LRPPRC, a multidomain organizer that potentially integrates mitochondria and the microtubular  $cytoskeleton\ with\ chromosome\ remodeling,\ for\ UXT\ increasing\ concentrations\ of\ UXT\ contributes\ to\ progressive$ aggregation of mitochondria and cell death potentially through its association with LRPPRC. UXT is possible a nuclear chaperone that promotes formation of the NF-kappa-B enhanceosome and which is essential for its nuclear function. It can suppresses cell transformation and mediate this function by interaction and inhibition of the biological activity of cell proliferation and survival stimulatory factors like MECOM

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Carrieri Claudia C	23064229	Nature	WB

Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 11047-1-AP (UXT antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



IP result of anti-UXT (IP:11047-1-AP, 3ug; Detection:11047-1-AP 1:300) with mouse brain tissue lysate 4000ug.