## For Research Use Only

# NME1 Polyclonal antibody

Catalog Number: 11086-2-AP

11 Publications



**Basic Information** 

Catalog Number:

11086-2-AP

Size:

550 µg/ml

Source:

Rabbit

Sotype:

GenBank Accession Number:

BC018994

GeneID (NCBI):

4830

UNIPROT ID:

P15531

Full Name:

IgG non-metastatic cells 1, protein (NM23A) expressed in

AG1548 Calculated MW:

17 kDa Observed MW: 16-20 kDa Purification Method: Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200 IF 1:20-1:200

**Applications** 

Tested Applications: IF/ICC, IHC, IP, WB, ELISA Cited Applications: WB, IF, IHC, CoIP

Species Specificity: human

**Cited Species:** 

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: HeLa cells, multi-cells

IP: A549 cells,

IHC: human lung cancer tissue, human pancreas

cancer tissue IF : MCF-7 cells,

# **Background Information**

NME1 is also named as NDPKA(Nucleoside diphosphate kinase A), NM23(Metastasis inhibition factor nm23), GAAD(Granzyme A-activated DNase) and belongs to the NDK family. It is a major role in the synthesis of nucleoside triphosphates other than ATP and required for neural development including neural patterning and cell fate determination. It has 3 isoforms produced by alternative splicing.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Anna Bianchi-Smiraglia	34667203	Nat Commun	WB
Marianna Naki	35563580	Int J Mol Sci	CoIP
Hongyuan Zhao	33786631	Oncol Rep	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




HeLa cells were subjected to SDS PAGE followed by western blot with 11086-2-AP (NME1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human lung cancer using 11086-2-AP (NME1 antibody) at dilution of 1:100 (under 10x lens)




Immunohistochemical analysis of paraffinembedded human lung cancer using 11086-2-AP (NME1 antibody) at dilution of 1:100 (under 40x lens).



WB result of 10086-2-AP (NME1 antibody).



IP result of anti-NME1 (IP:11086-2-AP, 4ug; Detection:11086-2-AP 1:500) with A549 cells lysate 2800ug.



Immunofluorescent analysis of MCF-7 cells using 11086-2-AP (NME1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).