For Research Use Only

FGFR4 Polyclonal antibody

Catalog Number: 11098-1-AP

Featured Product

18 Publications



Basic Information

Catalog Number: GenBank Accession Number: 11098-1-AP BC011847 GeneID (NCBI): Concentration: 600 ug/ml 2264 **UNIPROT ID:** Source: Rabbit P22455 Full Name: Isotype:

fibroblast growth factor receptor 4

Calculated MW: Immunogen Catalog Number: AG1573

88 kDa

Observed MW: 100-110 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:200-1:1000 IHC 1:50-1:500

IF/ICC 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IF Species Specificity:

human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HepG2 cells, mouse lung tissue IHC: human breast cancer tissue,

IF/ICC: HepG2 cells,

Background Information

Fibroblast growth factor receptor 4 (FGFR4) is a member of a highly conserved tyrosine kinase family, along with FGFR1-3. This family consists of an intracellular tyrosine kinase domain, a single transmembrane domain, and extracellular ligand binding domains (PMID:32492514). FGFR4 is the predominant FGFR isoform present in human $he patocytes. \, FGFR4 \, has \, been \, proposed \, to \, play \, a \, role \, in \, the \, observed \, induction \, of \, he patocyte \, proliferation \, and \, in \, the \, observed \, induction \, of \, he patocyte \, proliferation \, and \, in \, the \, observed \, induction \, of \, he patocyte \, proliferation \, and \, in \, the \, observed \, induction \, of \, he patocyte \, proliferation \, and \, in \, the \, observed \, i$ carcinogenesis by FGF19; however, contradicting evidence proposing a protective role for FGFR4 in suppressing hepatoma progression has also been proposed(PMID:20018895). While the role of FGFR4 in cancer remains to be fully elucidated, several findings suggest that this receptor may be an important player in Hepatocellular carcinoma (HCC) development and/or progression(PMID:10336501).

Notable Publications

Author	Pubmed ID	Journal	Application
Limin Wang	36305369	Tissue Eng Part A	IF
Yanan Chen	31628911	Biochem Pharmacol	WB
Yanwei Ye	32599983	Cancer Res Treat	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 11098-1-AP (FGFR4 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 11098-1-AP (FGFR4 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 11098-1-AP (FGFR4 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 11098-1-AP (FGFR4 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).