### For Research Use Only

# GATA2 Polyclonal antibody

Catalog Number: 11103-1-AP

Featured Product

15 Publications



**Basic Information** 

Catalog Number: 11103-1-AP Size:

650 μg/ml Source:

Rabbit
Isotype:
IgG
Immunogen Catalog Number:

AG1557

GenBank Accession Number:

BC018988 GeneID (NCBI): 2624

ENSEMBL Gene ID: ENSG00000179348 UNIPROT ID: P23769 Full Name:

GATA binding protein 2

Observed MW: 50-60 kDa Antigen affinity purification Recommended Dilutions: WB 1:500-1:3000 IHC 1:20-1:200 IF/ICC 1:50-1:500

**Purification Method:** 

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IF, ChIP

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Positive Controls:** 

WB: HeLa cells, HepG2 cells, mouse liver tissue, K-562 cells, mouse heart tissue, rat heart tissue, HEK-293

cells, rat liver tissue

IHC: human lymphoma tissue, IF/ICC: SH-SY5Y cells,

## **Background Information**

GATA2 is a member of GATA family of transcription factors that contains zinc fingers in DNA binding domain. GATA2 acts as a transcriptinal activator that regulates endothelin-1 expression in endothelial cells. Together with GATA3, GATA2 regulates adipocyte differentiation through molecular control of the preadpocyte-adipocyte transition. It also affect the activity of Rho inhibitor p190RhoGAP that controls capillary network formation in vitro and retinal angiogenesis in vivo.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Chao Shen	27617961	PLoS Genet	WB
Kazuya Kusama	34606582	Endocrinology	ChIP,IF
Joshua E Burda	35614216	Nature	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 11103-1-AP (GATA2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lymphoma using 11103-1-AP (GATA2 antibody) at dilution of 1:50 (under 10x lens).



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using GATA2 antibody (11103-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).