

For Research Use Only

GATA2 Polyclonal antibody

Catalog Number: 11103-1-AP

Featured Product

9 Publications



Basic Information

Catalog Number:

11103-1-AP

Size:

650 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1557

GenBank Accession Number:

BC018988

GeneID (NCBI):

2624

ENSEMBL Gene ID:

ENSG00000179348

UNIPROT ID:

P23769

Full Name:

GATA binding protein 2

Observed MW:

50-60 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IHC 1:20-1:200

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

WB, IF, IHC, ChIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, HepG2 cells, mouse liver tissue, K-562 cells, mouse heart tissue, rat heart tissue, HEK-293 cells, rat liver tissue

IHC : human lymphoma tissue,

Background Information

GATA2 is a member of GATA family of transcription factors that contains zinc fingers in DNA binding domain. GATA2 acts as a transcriptional activator that regulates endothelin-1 expression in endothelial cells. Together with GATA3, GATA2 regulates adipocyte differentiation through molecular control of the preadipocyte-adipocyte transition. It also affect the activity of Rho inhibitor p190RhoGAP that controls capillary network formation in vitro and retinal angiogenesis in vivo.

Notable Publications

Author	Pubmed ID	Journal	Application
Chao Shen	27617961	PLoS Genet	WB
Kazuya Kusama	34606582	Endocrinology	ChIP,IF
Joshua E Burda	35614216	Nature	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 11103-1-AP (GATA2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lymphoma using 11103-1-AP (GATA2 antibody) at dilution of 1:50 (under 10x lens).