

For Research Use Only

ACVR1 Polyclonal antibody

Catalog Number: 11108-1-AP



Basic Information

| | | |
|--|---|--|
| Catalog Number: 11108-1-AP | GenBank Accession Number: BC033867 | Purification Method: Antigen affinity purification |
| Size: 600 µg/ml | GeneID (NCBI): 90 | Recommended Dilutions: WB 1:500-1:1000 |
| Source: Rabbit | UNIPROT ID: Q04771 | |
| Isotype: IgG | Full Name: activin A receptor, type I | |
| Immunogen Catalog Number: AG1556 | Calculated MW: 509 aa, 57 kDa | |
| | Observed MW: 57 kDa | |

Applications

| | |
|--|--|
| Tested Applications: WB, ELISA | Positive Controls: WB : mouse kidney tissue, |
| Species Specificity: human, mouse, rat | |

Background Information

ACVR1 (activin receptor type I), also known as ALK2 or ACTRI, is a receptor for activin. It forms a stable complex with type II receptor after ligand binding. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling, and type II receptors are required for binding ligands and for expression of type I receptors. ACVR1 is expressed in many tissues including skeletal muscle and chondrocytes. It functions as a receptor for bone morphogenetic protein (BMP) and induces Indian hedgehog in chondrocytes during skeletal development. Mutations in ACVR1 gene are associated with fibrodysplasia ossificans progressive (PMID: 16642017).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 11108-1-AP (ACVR1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.