

For Research Use Only

# CYLD Polyclonal antibody

Catalog Number: 11110-1-AP

Featured Product

22 Publications



## Basic Information

**Catalog Number:**

11110-1-AP

**Size:**

550 µg/ml

**Source:**

Rabbit

**Isotype:**

IgG

**Immunogen Catalog Number:**

AG1598

**GenBank Accession Number:**

BC012342

**GeneID (NCBI):**

1540

**UNIPROT ID:**

Q9NQC7

**Full Name:**

cylindromatosis (turban tumor syndrome)

**Calculated MW:**

107 kDa

**Observed MW:**

110 kDa

**Purification Method:**

Antigen affinity purification

**Recommended Dilutions:**

WB 1:500-1:2000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

## Applications

**Tested Applications:**

IHC, IP, WB, ELISA

**Cited Applications:**

IF, IHC, IP, WB

**Species Specificity:**

human, mouse, rat

**Cited Species:**

human, rat, mouse

**Positive Controls:**

**WB:** mouse brain tissue, HEK-293 cells, A431 cells, Jurkat cells

**IP:** mouse brain tissue,

**IHC:** human colon cancer tissue, human brain tissue, human colon tissue

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

CYLD, also named as CYLD1, belongs to the peptidase C67 family. It is the protease that specifically cleaves 'Lys-63'-linked polyubiquitin chains. CYLD has endo-ubiquitinase activity and plays an important role in the regulation of pathways leading to NF-kappa-B activation. CYLD contributes to the regulation of cell survival, proliferation and differentiation via its effects on NF-kappa-B activation. It is a negative regulator of Wnt signaling. CYLD inhibits HDAC6 and thereby promotes acetylation of alpha-tubulin and stabilization of microtubules. CYLD plays a role in the regulation of microtubule dynamics, and thereby contributes to the regulation of cell proliferation, cell polarization, cell migration, and angiogenesis. It is required for normal cell cycle progress and normal cytokinesis. CYLD inhibits nuclear translocation of NF-kappa-B and plays a role in the regulation of inflammation and the innate immune response, via its effects on NF-kappa-B activation. It is dispensable for the maturation of intrathymic natural killer cells, but required for the continued survival of immature natural killer cells. CYLD negatively regulates TNFRSF11A signaling and osteoclastogenesis. This antibody is a rabbit polyclonal antibody raised against residues near the C terminus of human CYLD.

## Notable Publications

Author	Pubmed ID	Journal	Application
Hai-Yan Cui	34629821	World J Gastroenterol	WB
Xing Lin	27738385	Mediators Inflamm	WB
Guixin Zhu	34497368	Nat Cell Biol	WB

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 11110-1-AP (CYLD antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer using 11110-1-AP (CYLD antibody) at dilution of 1:50 (under 40x lens).



IP result of anti-CYLD (IP:11110-1-AP, 4ug; Detection:11110-1-AP 1:300) with mouse brain tissue lysate 4000ug.