For Research Use Only

MIRO2 Polyclonal antibody

Catalog Number: 11237-1-AP

Featured Product

26 Publications



Basic Information

Catalog Number: GenBank Accession Number: 11237-1-AP BC014942 GeneID (NCBI): Size: 700 ug/ml 89941 **UNIPROT ID:** Source:

Rabbit Q8IXI1 Isotype: Full Name:

IgG ras homolog gene family, member T2

Calculated MW: Immunogen Catalog Number:

AG1752 68 kDa

Observed MW: 80 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity: human, mouse **Cited Species:**

human, mouse, rat, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293T cells, human kidney tissue, mouse heart tissue, LO2 cells, HepG2 cells, human testis tissue, human heart tissue, K-562 cells, Hela cells

Purification Method:

WB 1:1000-1:10000

protein lysate

IHC 1:50-1:500

IF/ICC 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

IP: LO2 cells,

IHC: human colon tissue, human kidney tissue, human heart tissue, human lung cancer tissue

IF/ICC: HeLa cells,

Background Information

RHOT2 (Ras homolog gene family, member T2), also known as ARHT2, RASL or MIRO-2 (mitochondrial Rho GTPase 2), is a 618 amino acid single-pass type IV membrane protein as a member of the Rho family of GTPases. It is localized to the outer mitochondrial membrane and plays a role in mitochondrial trafficking and fusion-fission dynamics. RHOT2 is ubiquitously expressed with highest levels found in pancreas, heart, kidney and skeletal muscle. MIRO2 is highly phosphorylated at basal conditions, as evidenced by an immunoreactive band that electrophoretically migrates 7 kDa higher (~75 kDa) compared to its predicted molecular weight (68 kDa) (PMID: 28556983).

Notable Publications

Author	Pubmed ID	Journal	Application
Crystal A Lee	28924745	Protein Cell	WB,IF
Matthew Yoke Wui Ng	36270994	Nat Commun	WB
Tatiana Brustovetsky	34685760	Cells	WB, CoIP

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11237-1-AP (MIRO2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of MIRO 2 antibody (11237-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MIRO 2 transfected HepG2 cells.



IP result of anti-MIRO2 (IP:11237-1-AP, 4ug; Detection:11237-1-AP 1:4000) with LO2 cells lysate 3080 ug.



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 11237-1-AP (MIRO2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MIRO2 antibody (11237-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).