

For Research Use Only

PDI Polyclonal antibody

Catalog Number: 11245-1-AP

Featured Product

60 Publications



Basic Information

Catalog Number:

11245-1-AP

Size:

850 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1747

GenBank Accession Number:

BC014504

GeneID (NCBI):

5034

UNIPROT ID:

P07237

Full Name:

prolyl 4-hydroxylase, beta polypeptide

Calculated MW:

57 kDa

Observed MW:

57 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat, zebrafish

Cited Species:

human, mouse, rat, pig, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : COLO 320 cells, HEK-293 cells, HepG2 cells, mouse heart tissue, rat brain tissue

IP : HepG2 cells,

IHC : human prostate cancer tissue, human colon tissue

IF/ICC : HepG2 cells,

Background Information

PDIA1 (Protein disulfide-isomerase) is also named as ERBA2L, PDI, P4HB, PO4DB. It is a multifunctional protein that catalyzes the formation, breakage and rearrangement of disulfide bonds. In some cell types, it seems to be secreted or associated with the plasma membrane, where it undergoes constant shedding and replacement from intracellular sources. It can exist as homodimer and monomers and homotetramers may also occur (PMID:12095988).

Notable Publications

Author	Pubmed ID	Journal	Application
Liguo Yang	32943607	Cell Death Dis	IF
Linyu Sun	34555354	Mol Cell	IF
Nirav Florian Chhabra	34487921	Mol Metab	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11245-1-AP (PDI antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 11245-1-AP (PDI antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Anti-Human PDI (11245-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



WB result of PDI antibody (11245-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PDI transfected HEK-293 cells.



IP result of anti-PDI (IP:11245-1-AP, 4ug; Detection:11245-1-AP 1:2000) with HepG2 cells lysate 2600 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using PDI antibody (11245-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red). DAPI (blue).