For Research Use Only

RAD51AP1 Polyclonal antibody

Catalog Number:11255-1-AP

Featured Product

11 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com


Basic Information	Catalog Number: 11255-1-AP	GenBank Accession Number: BC016330	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	550 µg/ml	10635	WB 1:1000-1:4000	
	Source: Rabbit	UNIPROT ID: Q96B01	IF/ICC 1:50-1:500	
	Isotype:	Full Name:		
	IgG	RAD51 associated protein 1		
	Immunogen Catalog Number:	Calculated MW:		
	AG1781	38 kDa Observed MW:		
		38 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IF/ICC, ELISA	WB : HeLa cells, human brain tissue, mouse liver		
	Cited Applications: WB, IF, ChIP		tissue, rat liver tissue, Raji cells	
	Species Specificity: human, mouse, rat	IF/ICC : F	Hela cells,	
	Cited Species:			
	human, mouse			
Background Information	human, mouse Homologous recombination (HR) and other deleterious lesions from [PMID:12456786]. RAD51AP1 (RAI vertebrate-specific protein that er knockdown by siRNA leads to cell	n chromosomes. The eukaryotic HR re D51 associated protein 1), so named b nhances the ability of RAD51 to medi ular sensitivity to DNA damaging ag	eaction is mediated by RAD51 pecause it can interact with human RAD51, i ate the D-loop reaction. RAD51AP1 ents and elevated DNA damage-induced	
Background Information	human, mouse Homologous recombination (HR) and other deleterious lesions from [PMID:12456786]. RAD51AP1 (RAI vertebrate-specific protein that er knockdown by siRNA leads to cell	n chromosomes. The eukaryotic HR re D51 associated protein 1), so named b nhances the ability of RAD51 to medi- ular sensitivity to DNA damaging ag If the RAD51 recombinase activity by	eaction is mediated by RAD51 pecause it can interact with human RAD51, i ate the D-loop reaction. RAD51AP1 ents and elevated DNA damage-induced	
	human, mouse Homologous recombination (HR) and other deleterious lesions from [PMID:12456786]. RAD51AP1 (RAI vertebrate-specific protein that er knockdown by siRNA leads to cell chromatid breaks. Enhancement of	n chromosomes. The eukaryotic HR re D51 associated protein 1), so named b nhances the ability of RAD51 to medi- ular sensitivity to DNA damaging ag If the RAD51 recombinase activity by	eaction is mediated by RAD51 pecause it can interact with human RAD51, i ate the D-loop reaction. RAD51AP1 ents and elevated DNA damage-induced	
	human, mouse Homologous recombination (HR) i and other deleterious lesions from [PMID:12456786]. RAD51AP1 (RAI vertebrate-specific protein that er knockdown by siRNA leads to cell chromatid breaks. Enhancement c interaction of the two HR factors [I	n chromosomes. The eukaryotic HR re D51 associated protein 1), so named b nhances the ability of RAD51 to medi- ular sensitivity to DNA damaging ag of the RAD51 recombinase activity by PMID:17996710,17996711].	eaction is mediated by RAD51 because it can interact with human RAD51, i ate the D-loop reaction. RAD51AP1 ents and elevated DNA damage-induced RAD51AP1 is contingent upon the physical	
Background Information	human, mouse Homologous recombination (HR) i and other deleterious lesions from [PMID:12456786]. RAD51AP1 (RAI vertebrate-specific protein that er knockdown by siRNA leads to cell chromatid breaks. Enhancement c interaction of the two HR factors [I	n chromosomes. The eukaryotic HR re D51 associated protein 1), so named b hances the ability of RAD51 to medi- ular sensitivity to DNA damaging ag of the RAD51 recombinase activity by PMID:17996710,17996711]. Pubmed ID Journal	pecause it can interact with human RAD51, is ate the D-loop reaction. RAD51AP1 ents and elevated DNA damage-induced RAD51AP1 is contingent upon the physical Application	
	human, mouse Homologous recombination (HR) i and other deleterious lesions from [PMID:12456786]. RAD51AP1 (RAI vertebrate-specific protein that er knockdown by siRNA leads to cell chromatid breaks. Enhancement co interaction of the two HR factors [I Author Tribhuwan Yadav	n chromosomes. The eukaryotic HR re D51 associated protein 1), so named b nhances the ability of RAD51 to medi- ular sensitivity to DNA damaging ag of the RAD51 recombinase activity by PMID:17996710,17996711]. Pubmed ID Journal 36265486 Mol Cell	eaction is mediated by RAD51 because it can interact with human RAD51, i ate the D-loop reaction. RAD51AP1 ents and elevated DNA damage-induced RAD51AP1 is contingent upon the physical Application WB	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 11255-1-AP (RAD51AP1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 11255-1-AP (RAD51AP1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).