For Research Use Only

EIF3H Polyclonal antibody

Catalog Number:11310-1-AP

Featured Product

5 Publications

BC000386

8667

015372 Full Name:

GeneID (NCBI):

UNIPROT ID:

factor 3, subunit H

Calculated MW:

GenBank Accession Number:

eukaryotic translation initiation



Basic Information

Catalog Number: 11310-1-AP Size:

550 μg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG1850

40 kDa Observed MW:

40 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:3000

Positive Controls:

WB: HeLa cells, NIH/3T3 cells, PC-12 cells

Applications

Tested Applications: WB, ELISA

Cited Applications:

WB, IP, IF

Species Specificity: human, mouse, rat Cited Species: human

Background Information

Eukaryotic translation initiation factor eIF3, that plays a central role in translation initiation, consists of five core subunits that are present in both the budding yeast and higher eukaryotes. The EIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex. The complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. It is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. EIF3H is a component of the EIF3 complex. [PMID:19204574,20503360]

Notable Publications

| Author | Pubmed ID | Journal | Application |
|------------|-----------|---------------------------|-------------|
| Yuanpei Li | 36289222 | Nat Commun | WB |
| Genhua Yu | 30022709 | Biosci Biotechnol Biochem | WB |
| Mengqi Chu | 33509274 | Stem Cell Res Ther | WB |

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 11310-1-AP (EIF3H antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.