For Research Use Only

NASP Polyclonal antibody

Catalog Number: 11323-1-AP

Featured Product

11 Publications



Basic Information

Catalog Number: GenBank Accession Number: 11323-1-AP BC010105 GeneID (NCBI): Size: 700 μg/ml 4678 **UNIPROT ID:** Source: Rabbit P49321

Full Name: Isotype: nuclear autoantigenic sperm protein (histone-binding) Immunogen Catalog Number:

AG1824 Calculated MW: 788 aa. 85 kDa Observed MW: 150 kDa, 70 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA **Cited Applications:**

WB, IHC, CoIP Species Specificity: human, mouse, rat **Cited Species:** human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse testis tissue, IP: mouse testis tissue,

IHC: human testis tissue, human lymphoma tissue

Purification Method:

WB 1:500-1:2400

protein lysate

IHC 1:20-1:200

IF/ICC 1:10-1:100

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

IF/ICC: MCF-7 cells,

Background Information

 $NASP \ (nuclear \ autoantigenic \ sperm \ protein) \ is \ associated \ with \ DNA \ replication, cell \ proliferation \ and \ cell \ cycle$ progression through functioning as a Histone H1 binding protein that mediates histone transport to the nucleus. NASP had two isoforms, tNASP (testis type) and sNASP (somatic type). The tNASP was mainly expressed in testis, a variety of malignant tumors, stem cells and embryonic tissues, while sNASP existed in all somatic mitosis cells. This antibody can recognize both isoforms.

Notable Publications

Author	Pubmed ID	Journal	Application
Fei Xiong	36167790	Cell Death Dis	WB
Qing Yuan	25378924	Int J Nanomedicine	WB
Nishibu Takahiro T	23229118	Biosci Trends	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




mouse testis tissue were subjected to SDS PAGE followed by western blot with 11323-1-AP (NASP antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human testis using 11323-1-AP (NASP antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human testis using 11323-1-AP (NASP antibody) at dilution of 1:100 (under 40x lens).



Immun of luorescent analysis of MCF-7 cells, using NASP antibody 11323-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-NASP (IP:11323-1-AP, 5ug; Detection:11323-1-AP 1:1000) with mouse testis tissue lysate 4000ug.