For Research Use Only


Integrin Beta 7 Polyclonal antibody Catalog Number:11328-1-AP 6 Publications




Basic Information	Catalog Number: 11328-1-AP	GenBank Accession Nu BC015916	ımber:	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):		Recommended Dilutions:
	600 µg/ml	3695		WB 1:200-1:1000
	Source: Rabbit	UNIPROT ID: P26010		IHC 1:50-1:500
	lsotype: IgG	Full Name: integrin, beta 7		
	Immunogen Catalog Number: AG1871	Calculated MW: 798 aa, 87 kDa		
		Observed MW: 87-100 kDa		
Applications	Tested Applications: WB, IHC, ELISA		Positive Cor	
	Cited Applications: WB: THP-1 c WB, IHC, IF WB: the three thr		ells, human lung tissue, K-562 cells, en tissue	
	Species Specificity: human, mouse, rat		IHC : human	tonsillitis tissue,
	Cited Species: human, mouse, rat			
	Note-IHC: suggested antig TE buffer pH 9.0; (*) Alter retrieval may be performe buffer pH 6.0	natively, antigen		
Background Information	ITGB7 (integrin, beta7) is a single-pass type I membrane protein that belongs to the integrin beta chain family. The integrins are a large family of cell-surface glycoprotein receptors that play key roles in leucocyte adhesion, signalling, proliferation and migration by binding as heterodimers to specific ligands. The beta7 integrin associates with alpha4 integrin to form a heterodimer, alpha4/beta7, which interacts with MADCAM1, VCAM, fibronectin, and may also interact with HIV-1 gp120. The beta7 integrin can also form a heterodimer with alphaE integrin, is a receptor for E-cadherin. ITGB7 is expressed in a variety of leukocyte lines.			
Notable Publications	Author	Pubmed ID Journal	1	Application
	Tong Sun		mmun Signal	
	Gilad Halpert	24764299 J Biol C	•	IF
	Qi-Cheng Ni		pharmacol	WB
Storage	Storage: Store at -20°C. Stable for one yes Storage Buffer: PBS with 0.02% sodium azide a			


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





THP-1 cells were subjected to SDS PAGE followed by western blot with 11328-1-AP (Integrin Beta 7 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 11328-1-AP (Integrin Beta 7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 11328-1-AP (Integrin Beta 7 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).