

For Research Use Only

PSMC3IP Polyclonal antibody

Catalog Number: 11339-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number:

11339-1-AP

Size:

700 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1883

GenBank Accession Number:

BC008792

GeneID (NCBI):

29893

UNIPROT ID:

Q9P2W1

Full Name:

PSMC3 interacting protein

Calculated MW:

25 kDa

Observed MW:

24-29 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:400-1:1600

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HEK-293 cells, K-562 cells, Jurkat cells

IP : Jurkat cells,

IHC : mouse testis tissue, rat testis tissue

IF/ICC : HepG2 cells,

Background Information

PSMC3IP, also called HOP2, encodes a protein that functions in meiotic recombination. It is a subunit of the PSMC3IP/MND1 complex, which interacts with PSMC3/TBP1 to stimulate DMC1- and RAD51-mediated strand exchange during meiosis. The protein encoded by this gene can also co-activate ligand-driven transcription mediated by estrogen, androgen, glucocorticoid, progesterone, and thyroid nuclear receptors. Mutations in this gene cause XX female gonadal dysgenesis.

Notable Publications

Author	Pubmed ID	Journal	Application
Nam Woo Cho	25259924	Cell	WB
Chih-Ying Lee	30622262	Nat Commun	
Matthew Tsang	30701021	Oncotarget	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11339-1-AP (HOP2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 11339-1-AP (HOP2 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 11339-1-AP (HOP2 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-HOP2 (IP:11339-1-AP, 4ug; Detection:11339-1-AP 1:500) with Jurkat cells lysate 3200ug.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using HOP2 antibody (11339-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).