#### For Research Use Only

# ZBTB7B Polyclonal antibody

Catalog Number: 11341-1-AP

4 Publications



**Basic Information** 

Catalog Number: 11341-1-AP

Size: 800  $\mu$  g/ml Source:

Rabbit Isotype: IgG

Immunogen Catalog Number: AG1888

GenBank Accession Number:

BC012070 GeneID (NCBI): 51043 Full Name:

zinc finger and BTB domain containing 7B

Calculated MW: 539 aa, 58 kDa Observed MW: 60-65 kDa

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:4000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF/ICC 1:800-1:3200

**Applications** 

**Tested Applications:** WB, IF/ICC, IP, ELISA Cited Applications:

WB, IHC, IP Species Specificity: human, mouse, rat Cited Species:

human, mouse, bovine, hamster

Positive Controls:

WB: HeLa cells, HepG2 cells

IP: HeLa cells, IF/ICC: HeLa cells,

### **Background Information**

ZBTB7B belongs to a large family of transcription factors, generally acting as repressors, characterized by a carboxyterminal DNA binding domain made of multiple zinc fingers (four in Thpok) and an amino-terminal BTB-POZ domain that mediates homo- (and possibly hetero-) dimerization [PMID: 17084908]. Zbtb7b is up-regulated by MHC-II-restricted thymocytes during their CD4 differentiation and is a major determinant of CD4 lineage choice. Two properties of ZBTB7B deserve attention. First, although Thpok is expressed in a wide variety of cells, its expression in the thymus is highly lineage-specific: CD4 SP thymocytes (and all CD4 T cells) express Thpok, whereas DP and CD8 SP thymocytes do notSecond, both loss- and gain-of-function experiments indicate that Thpok affects lineage choice but not positive selection [PMID: 15729333, 10550051].

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jian Sun	34700042	Acta Biomater	WB
Hao-Ming Xu	35761286	J Transl Med	WB,IHC
Jiaxin Wang	38154091	J Agric Food Chem	WB,IP

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 11341-1-AP (ZBTB7B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



IP result of anti-ZBTB7B (IP:11341-1-AP, 4ug; Detection:11341-1-AP 1:3000) with HeLa cells lysate 1320 ug.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using ZBTB7B antibody (11341-1-AP) at dilution of 1:1600 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).