For Research Use Only

SPRY2 Polyclonal antibody

Catalog Number: 11383-1-AP

Featured Product

8 Publications



Basic Information

Catalog Number: GenBank Accession Number: 11383-1-AP BC015745
Size: GeneID (NCBI): 10253
Source: UNIPROT ID: Rabbit O43597
Isotype: Full Name:

sprouty homolog 2 (Drosophila)

Immunogen Catalog Number:Calculated MW:AG191835 kDaObserved MW:

35-40 kDa, 60~65 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IF Species Specificity:

human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: SW 1990 cells, Jurkat cells, A375 cells, Neuro-2a

Purification Method:

WB 1:500-1:1000 IHC 1:50-1:500

IF/ICC 1:10-1:100

Antigen affinity purification

Recommended Dilutions:

cells

IHC: human pancreas cancer tissue,

IF/ICC: A375 cells,

Background Information

Sprouty(SPRY) genes encode intracellular inhibitors of receptor tyrosine kinase (RTK) signaling pathways, including those triggered by fibroblast growth factors (Fgfs). In addition to its well-reported function as an inhibitor of the RAS/RAF/MEK signalling cascade, SPRY2 exerts its inhibitory role in the PI3K/AKT signalling pathway through PTEN, and inhibits ERK/MAPK activity via EGFR trafficking. SPRY2 was detected 60-65 kDa (PMID:29291435).

Notable Publications

Author	Pubmed ID	Journal	Application
Takaharu Taketomi	28921936	Cell Biol Int	WB
Guangying Yang	35682694	Int J Mol Sci	WB
Nai-Jie Fu	35307375	Toxicol Appl Pharmacol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11383-1-AP (SPRY2 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.




SW 1990 cells were subjected to SDS PAGE followed by western blot with 11383-1-AP (SPRY2 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 11383-1-AP (SPRY2 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 11383-1-AP (SPRY2 Antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of A375 cells using 11383-1-AP (SPRY2 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated Goat Anti-Rabbit lgG(H+L).



A375 cells were subjected to SDS PAGE followed by western blot with 11383-1-AP (SPRY2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.