

For Research Use Only

PSMC4 Polyclonal antibody

Catalog Number: 11389-1-AP **5 Publications**



Basic Information

Catalog Number: 11389-1-AP	GenBank Accession Number: BC014488	Purification Method: Antigen affinity purification
Size: 550 µg/ml	GeneID (NCBI): 5704	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500 IF 1:10-1:100
Source: Rabbit	UNIPROT ID: P43686	
Isotype: IgG	Full Name: proteasome (prosome, macropain) 26S subunit, ATPase, 4	
Immunogen Catalog Number: AG1943	Calculated MW: 418 aa, 47 kDa	
	Observed MW: 43-50 kDa	

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA	Positive Controls: WB : mouse brain tissue, rat brain tissue IHC : human colon cancer tissue, IF : HepG2 cells, SH-SY5Y cells
Cited Applications: WB, IF, IHC	
Species Specificity: human, mouse, rat	
Cited Species: human, rat, mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

PSMC4 (proteasome (prosome, macropain) 26S subunit, ATPase, 4) is also named as TBP7, S6, TBP7, MIP224 and belongs to the AAA ATPase family. The 26S protease is involved in the ATP-dependent degradation of ubiquitinated proteins. The predicted 458-amino acid TBP7 protein is 57% identical to TBP1 and they can form heterodimers (PMID:8419915).

Notable Publications

Author	Pubmed ID	Journal	Application
Indrajit Sahu	34702852	Nat Commun	WB
Ling Lin	34836374	Nutrients	WB
Lien Nguyen	31831332	Neuron	IF

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 11389-1-AP (PSMC4 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 11389-1-AP (PSMC4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells, using PSMC4 antibody 11389-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).