For Research Use Only

NMNAT1 Polyclonal antibody

Catalog Number: 11399-1-AP

Featured Product

10 Publications



Purification Method:

WB 1:500-1:1000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 11399-1-AP BC014943 GeneID (NCBI): Concentration: 500 ug/ml 64802 **UNIPROT ID:** Source: Rabbit Q9HAN9 Full Name: Isotype:

nicotinamide nucleotide adenylyltransferase 1 Immunogen Catalog Number:

AG1944 Calculated MW: 33 kDa

Observed MW: 28-32 kDa

Applications

Tested Applications: WB, IHC, ELISA **Cited Applications:** WB, IHC, IF

Species Specificity: human **Cited Species:** human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human testis tissue.

IHC: human pancreas cancer tissue,

Background Information

 $NMNAT1 is a \ member of the \ nicotina mide-nucleotide \ adenylyl transferases (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ adenylyl transferases (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ adenylyl transferases (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ adenylyl transferases (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ adenylyl transferases (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ adenylyl transferases (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ adenylyl transferases (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ adenylyl transferases \ (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ adenylyl transferases \ (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ adenylyl transferases \ (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ adenylyl transferases \ (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ adenylyl transferases \ (NMNATs) \ which \ catalyze \ nicotina mide-nucleotide \ nic$ adenine dinucleotide (NAD) synthesis (PMID: 28445802). NMNAT is a central enzyme in NAD biosynthesis, catalyzing the formation of NAD+ from nicotinamide mononucleotide (NMN) and ATP (PMID: 17402747). NMNAT1 is widely expressed with the highest levels in skeletal muscle, heart, and kidney (PMID: 11027696). Mutations in NMNAT1 have been shown associated with the LCA9 form of the retinal degeneration pathology Leber's congenital amaurosis (PMID: 22842229, 22842230).

Notable Publications

Author	Pubmed ID	Journal	Application
Ya Li	33053398	Metabolism	IF
Yan Liu	33793246	J Proteome Res	WB
Itaru Yasuda	33795425	J Am Soc Nephrol	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human testis tissue were subjected to SDS PAGE followed by western blot with 11399-1-AP (NMNAT1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 11399-1-AP (NMNAT1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 11399-1-AP (NMNAT1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).