For Research Use Only

PRAME Polyclonal antibody

Catalog Number: 11438-1-AP



Purification Method:

WB 1:500-1:1000

protein lysate

IHC 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 11438-1-AP BC014074 GeneID (NCBI): Size: 450 µg/ml 23532 **UNIPROT ID:** Source: Rabbit P78395

Full Name: Isotype: preferentially expressed antigen in

melanoma

50-58 kDa

AG1906 Calculated MW: 509 aa, 58 kDa Observed MW:

Immunogen Catalog Number:

Applications

Tested Applications: WB, IP, IHC, ELISA WB: K-562 cells, Species Specificity: IP: K-562 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human lung cancer tissue,

Background Information

The PRAME (preferentially expressed antigen of melanoma) gene was previously shown to be overexpressed in ovarian/primary peritoneal serous carcinoma compared with malignant mesothelioma using gene expression arrays. It is considered a melanocyte differentiation antigen which is overexpressed in both solid and hematologic tumors. In normal tissue, a very low level of PRAME expression is found in normal testis, adrenals, ovary and endometrium. A high level of PRAME expression has been reported for several solid tumors, including ovarian cancer, breast cancer, lung cancer and melanomas, medulloblastoma, sarcomas, head and neck cancers, neuroblastoma, renal cancer, and Wilms'tumor. As a nuclear transcriptional repressor protein, PRAME binds to retinoic acid receptor a, thereby inhibiting retinoic acid induced differentiation, growth arrest, and apoptosis.


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



K-562 cells were subjected to SDS PAGE followed by western blot with 11438-1-AP (PRAME antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-PRAME (IP:11438-1-AP, 4ug; Detection:11438-1-AP 1:300) with K-562 cells lysate 2400 ug.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 11438-1-AP (PRAME antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).