

For Research Use Only

COPE Polyclonal antibody

Catalog Number: 11457-1-AP



Basic Information

Catalog Number:

11457-1-AP

Size:

133 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG2024

GenBank Accession Number:

BC017285

GeneID (NCBI):

11316

UNIPROT ID:

O14579

Full Name:

coatamer protein complex, subunit

epsilon

Calculated MW:

34 kDa

Observed MW:

30-35 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IHC 1:50-1:500

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HEK-293 cells, PC-3 cells, A375 cells, HeLa cells, A431 cells, MCF-7 cells

IHC : human stomach cancer tissue, human liver cancer tissue, human prostate cancer tissue, rat stomach tissue

IF/ICC : HepG2 cells,

Background Information

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HEK-293 cells were subjected to SDS PAGE followed by western blot with 11457-1-AP (COPE antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 11457-1-AP (COPE antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 11457-1-AP (COPE antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using COPE antibody (11457-1-AP) at dilution of 1:400 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).