For Research Use Only

GPSM1 Polyclonal antibody

Catalog Number: 11483-1-AP

Featured Product

4 Publications



Basic Information

Catalog Number:

11483-1-AP

BC017353

Source:

Rabbit

26086

Isotype:

IgG

UNIPROT ID:

Q86YR5

Immunogen Catalog Number:

Full Name:

Full Name: G-protein signaling modulator 1

Calculated MW: 652 aa, 72 kDa Observed MW: 70-80 kDa

(AGS3-like, C. elegans)

Purification Method: Antigen affinity purification Recommended Dilutions: WB: 1:500-1:3000

IHC: 1:50-1:500

IF/ICC: 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IF Species Specificity: human, mouse, rat Cited Species:

mouse, rat

AG2018

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 .

IF/ICC: HepG2 cells,

Positive Controls:

WB: mouse brain tissue, HeLa cells, human brain tissue, rat brain tissue

 $\hbox{IHC:} \hbox{ human oesophagus cancer tissue, mouse brain}\\$

tissue

Background Information

GPSM1, also named as AGS3, is a receptor-independent G protein activator that has been implicated in multiple biological events such as brain development, neuroplasticity and addiction, cardiac function, Golgi structure/function, macroautophagy and metabolism. It contains seven tetratricopeptide repeats in its N-terminal half and four G-protein regulatory (GPR) motifs in its C-terminal half. It has been shown that AGS3 could regulate the orientation of the mitotic spindle, cAMP production, membrane protein transport, and asymmetric cell division by binding preferentially to inactive Gai/o subunits complexed with guanine dinucleotide phosphate (GDP) at multiple G-protein regulatory or Goloco motif repeats. It also plays an important anti-apoptotic role through enhancing the phosphorylation of cyclic AMP response element-binding protein (p-CREB).

Notable Publications

Application
Application
WB,IHC,IF
WB.IHC
WB,IF

Storage

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 11483-1-AP (AGS3 antibody) at dilution of 1:600 incubated at room temperature for 15 bours




Various lysates were subjected to SDS PAGE followed by western blot with 11483-1-AP (GPSM1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 11483-1-AP (GPSM1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 11483-1-AP (GPSM1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human oesophagus cancer tissue slide using 11483-1-AP (GPSM1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human oesophagus cancer tissue slide using 11483-1-AP (GPSM1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using GPSM1 antibody (11483-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).