## For Research Use Only

# TFDP2 Polyclonal antibody

Catalog Number:11500-1-AP 3 Publications



**Basic Information** 

Catalog Number: 11500-1-AP Concentration: 700 ug/ml

Rabbit Q14188 Full Name: Isotype:

Immunogen Catalog Number:

AG2054

Source:

GenBank Accession Number:

BC021113 GeneID (NCBI): 7029 **UNIPROT ID:** 

transcription factor Dp-2 (E2F dimerization partner 2)

Calculated MW: 446 aa, 49 kDa Observed MW: 49 kDa

**Purification Method:** Antigen affinity purification Recommended Dilutions:

WB: 1:500-1:1000 IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

**Applications** 

**Tested Applications:** WB, IP, ELISA

Cited Applications:

WB, IHC

Species Specificity: human, mouse Cited Species: human, mouse

Positive Controls:

WB: mouse heart tissue, human brain tissue

IP: SH-SY5Y cells,

# **Background Information**

TFDP2, also known as E2FDP2, is a coactivator of E2F factors and involves in EGF or IFN-gamma-mediated effects by enhancing or reducing active heterodimeric complexes. The DP2/E2F complex functions in the control of cell-cycle progression from G1 to S phase. The E2F1/DP complex appears to mediate both cell proliferation and apoptosis [PMID:16360038]. Binds DNA cooperatively with E2F family members through the E2 recognition site, 5'-TTTC[CG]CGC-3', found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication [PMID:16721044].

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Caron Etienne E	21952136	Mol Syst Biol	WB
Shuwei Zhang	39793695	Neuropharmacology	WB
Ya-Jun Deng	38187091	Front Mol Biosci	IHC


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




mouse heart tissue were subjected to SDS PAGE followed by western blot with 11500-1-AP (TFDP2 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



human brain tissue were subjected to SDS PAGE followed by western blot with 11500-1-AP (TFDP2 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



IP result of anti-TFDP2 (IP:11500-1-AP, 4ug; Detection:11500-1-AP 1:400) with SH-SY5Y cells lysate 1600ug.



IP result of anti-TFDP2 (IP:11500-1-AP, 4ug; Detection:11500-1-AP 1:600) with SH-SY5Y cells lysate 1360 ug.