For Research Use Only

## TFDP2 Polyclonal antibody

Catalog Number:11500-1-AP

1 Publications





Basic Information	Catalog Number: GenBank Accession Number: 11500-1-AP BC021113		ession Number:	Purification Method: Antigen affinity purification	
	Size:	Genel D (NCB	):	Recommended Dilutions: WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	700 µg/ml	7029			
	Source: Rabbit	UNIPROT ID: Q14188 Full Name: transcription factor Dp-2 (E2F dimerization partner 2)		protein lysate	
	Isotype: IgG Immunogen Catalog Number:				
	AG2054	Calculated MW: 446 aa, 49 kDa			
		Observed MW 49 kDa	Observed MW: 49 kDa		
Applications	Tested Applications: IP, WB, ELISA		Positive Controls:		
	Cited Applications: WB	Cited Applications: IP : SH-SY5Y cells.			
	Species Specificity: human, mouse				
	Cited Species: mouse				
Background Information	TFDP2, also known as E2FDP2, is a coactivator of E2F factors and involves in EGF or IFN-gamma-mediated effects enhancing or reducing active heterodimeric complexes. The DP2/E2F complex functions in the control of cell-cycle progression from G1 to S phase. The E2F1/DP complex appears to mediate both cell proliferation and apoptosis [PMID:16360038]. Binds DNA cooperatively with E2F family members through the E2 recognition site, 5'- TTTC[CG]CGC-3', found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication [PMID:16721044].				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Caron Etienne E	21952136	Mol Syst Biol	WB	
Storage	Storage: Store at -20°C. Stable for one yes Storage Buffer:	ar after shipment.			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data





mouse heart tissue were subjected to SDS PAGE followed by western blot with 11500-1-AP (TFDP2 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.

human brain tissue were subjected to SDS PAGE followed by western blot with 11500-1-AP (TFDP2 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



IP result of anti-TFDP2 (IP:11500-1-AP, 4ug: Detection:11500-1-AP 1:400) with SH-SY5Y cells lysate 1600ug.