

For Research Use Only

ASGR2 Polyclonal antibody

Catalog Number: 11501-2-AP



Basic Information

Catalog Number: 11501-2-AP	GenBank Accession Number: BC017251	Purification Method: Antigen affinity purification
Size: 550 µg/ml	GeneID (NCBI): 433	Recommended Dilutions: WB 1:200-1:1000
Source: Rabbit	UNIPROT ID: P07307	IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate
Isotype: IgG	Full Name: asialoglycoprotein receptor 2	IHC 1:50-1:500
Immunogen Catalog Number: AG2037	Calculated MW: 39 kDa	
	Observed MW: 39 kDa	

Applications

Tested Applications:
FC, IHC, IP, WB, ELISA

Species Specificity:
human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : human liver tissue,

IP : L02 cells,

IHC : human liver tissue,

Background Information

Asialoglycoprotein receptor (ASGPR), also known as the hepatic galactose/N-acetylglucosamine (GlcNAc) receptor or Ashwell receptor, is a C-type lectin expressed exclusively in hepatic parenchymal cells. ASGPR consists of two subunits, a major subunit (ASGR1, HL-1) and a minor subunit (ASGR2, HL-2), and specifically recognizes terminal β -linked galactose or GlcNAc on circulating glycoproteins or cells. This receptor plays a critical role in serum glycoprotein homeostasis by mediating the endocytosis and lysosomal degradation of glycoproteins that contain terminal galactose or GlcNAc residues. ASGPR may facilitate hepatic infection by multiple viruses including hepatitis B, and is also a target for liver-specific drug delivery.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human liver tissue were subjected to SDS PAGE followed by western blot with 11501-2-AP (ASGR2 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.




IP result of anti-ASGR2 (IP:11501-2-AP, 4ug; Detection:11501-2-AP 1:300) with L02 cells lysate 1800ug.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 11501-2-AP (ASGR2 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 11501-2-AP (ASGR2 Antibody) at dilution of 1:200 (under 40x lens).



1x10⁶ HepG2 cells were stained with 0.2ug ASGR2 antibody (11501-2-AP, red) and control antibody (blue). Fixed with 4% PFA blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1500.