For Research Use Only

MOXD1 Polyclonal antibody

Catalog Number: 11545-1-AP

2 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

11545-1-AP

BC018756

Size:

GeneID (NCBI):

26002

Source:

UNIPROT ID:

Rabbit

Q6UVY6

Isotype:

Full Name:

monooxygenase, DBH-like 1

Immunogen Catalog Number:Calculated MW:AG206069 kDa

Observed MW: 63-70 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500

IF-P 1:50-1:500

Applications

Tested Applications: WB, IHC, IF-P, ELISA Cited Applications: WB, IHC

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, rat brain tissue IHC: mouse brain tissue, rat brain tissue

IF-P: mouse brain tissue,

Background Information

Monooxygenase DBH like 1 (MOXD1) is a member of the copper monooxygenase family, including dopamine monooxygenase (dopamine β -monooxygenase, DBM) and peptidylglycine α -hydroxylating monooxygenase (PHM). MOXD1 is expressed during the development of the neural crest, and distributed along the olfactory bulb, cerebellum, brain stem, and parietal cortex with the development of nerve cells, participating in the basal plate. MOXD1 is highly expressed in GBM cells and can promote the growth of cancer cells (PMID: 35393406). In addition, MOXD1 is a gate-keeper of organ homeostasis and functions as a tumor-suppressor in neuroblastoma

Notable Publications

Author	Pubmed ID	Journal	Application
Tongtong Pan	38042280	Life Sci	WB,IHC
Fuhai Bai	37889347	Funct Integr Genomics	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11545-1-AP (MOXD1 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 11545-1-AP (MOXD1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 11545-1-AP (MOXD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 11545-1-AP (MOXD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 11545-1-AP (MOXD1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using MOXD1 antibody (11545-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using MOXD1 antibody (11545-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).