For Research Use Only

DGKA Polyclonal antibody

Catalog Number: 11547-1-AP

Featured Product

19 Publications

BC023523

1606

P23743

GeneID (NCBI):

UNIPROT ID:

Full Name:

GenBank Accession Number:

diacylglycerol kinase, alpha 80kDa



Basic Information

Catalog Number: 11547-1-AP Size:

300 μg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG2075

Calculated MW: 735 aa, 83 kDa

> Observed MW: 83 kDa

> > Positive Controls:

WB: Jurkat cells, HeLa cells, MOLT-4 cells, human cerebellum tissue, HEK-293 cells, mouse cerebellum

Purification Method:

WB 1:2000-1:12000

protein lysate

IHC 1:50-1:500

IF 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

tissue IP: Jurkat cells,

IHC: human gliomas tissue, human brain tissue

IF: Jurkat cells,

Applications

Tested Applications:

FC, IF/ICC, IHC, IP, WB, ELISA

Cited Applications: WB, IP, IF, IHC Species Specificity: human, mouse, rat **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

DGKA(Diacylglycerol kinase alpha) is also named as DAGK, DAGK1 and belongs to the eukaryotic diacylglycerol kinase family, which is suggested to attenuate diacylglycerol-induced cell responses through the phosphorylation of this second messenger to phosphatidic acid.DGKA regulates the secretion of lethal exosomes bearing Fas ligand during activation-induced cell death of T lymphocytes.

Notable Publications

Author	Pubmed ID	Journal	Application
Naoki Okada	34482409	Cancer Immunol Immunother	IHC
Yi Kong	27498782	Oncol Rep	IHC
David Novo	30498210	Nat Commun	

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11547-1-AP (DGKA antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




IP result of anti-DGKA (IP:11547-1-AP, 3ug; Detection:11547-1-AP 1:1000) with Jurkat cells lysate 1500ug.




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 11547-1-AP (DGKA antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 11547-1-AP (DGKA antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using DGKA antibody (11547-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human DGKA (11547-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).