For Research Use Only

SRP68 Polyclonal antibody

Catalog Number: 11585-1-AP

Featured Product

9 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 11585-1-AP
 BC020238

 Size:
 GeneID (NCBI):

 800 ug/ml
 6730

 Source:
 UNIPROT ID:

 Rabbit
 Q9UHB9

 Isotype:
 Full Name:

Isotype: Full Name:
IgG signal recognition particle 68kDa

Immunogen Catalog Number: Calculated MW:

AG2177 68 kDa

Observed MW: 68 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000

IF/ICC 1:200-1:800

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species:

buffer pH 6.0

human, mouse

Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate

Positive Controls:

WB: mouse brain tissue, human brain tissue

IP: mouse kidney tissue,

IHC: human gliomas tissue, rat testis tissue

IF/ICC: HepG2 cells,

Background Information

The signal recognition particle (SRP) is a ribonucleoprotein complex that mediates the targeting of secretory proteins to the rough endoplasmic reticulum (ER) membrane. This ribonucleoprotein complex might interact directly with the docking protein in the ER membrane and possibly participate in the elongation arrest function. SRP68 is one component of the ribonucleprotein complex, and forms a heterodimer with SRP72.

Notable Publications

Author	Pubmed ID	Journal	Application
Mehdi Pirouz	32457326	Nat Commun	WB
Joseph Russo	28129347	PLoS One	WB
Piazzon Nathalie N	23221635	Nucleic Acids Res	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




mouse brain tissue were subjected to SDS PAGE followed by western blot with 11585-1-AP (SRP68 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 11585-1-AP (SRP68 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat testis tissue slide using 11585-1-AP (SRP68 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat testis tissue slide using 11585-1-AP (SRP68 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-SRP68 (IP:11585-1-AP, 4ug; Detection:11585-1-AP 1:400) with mouse kidney tissue lysate 4000ug.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using SRP68 antibody (11585-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).