

For Research Use Only

PPIH Polyclonal antibody

Catalog Number: 11651-1-AP

1 Publications



Basic Information

Catalog Number:

11651-1-AP

Concentration:

450 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG2253

GenBank Accession Number:

BC003412

GeneID (NCBI):

10465

UNIPROT ID:

O43447

Full Name:

peptidylprolyl isomerase H
(cyclophilin H)

Calculated MW:

19 kDa

Observed MW:

19 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:200-1:1000

IHC: 1:100-1:400

IF/ICC: 1:10-1:100

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

IF

Species Specificity:

human, mouse, rat

Cited Species:

human

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB: U2OS cells, K-562 cells, PC-3 cells, mouse colon
tissue, rat colon tissue

IHC: human colon cancer tissue,

IF/ICC: PC-3 cells,

Background Information

PPIH (Peptidyl-prolyl cis-trans isomerase H) is also named as CYP20, CYPH and belongs to cyclophilin-type PPIase family, PPIase H subfamily. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may act as a chaperone (PMID:12875835).

Notable Publications

Author	Pubmed ID	Journal	Application
Yinan Chen	34976220	Theranostics	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




U2OS cells were subjected to SDS PAGE followed by western blot with 11651-1-AP (PPIH antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 11651-1-AP (PPIH antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 11651-1-AP (PPIH Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 11651-1-AP (PPIH Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of PC-3 cells using 11651-1-AP (PPIH antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).